

STRINGY POWER OPERATIONS IN TATE K-THEORY

NORA GANTER

ABSTRACT. We study the loop spaces of the symmetric powers of an orbifold and use our results to define equivariant power operations in Tate K-theory. We prove that these power operations are elliptic and that the Witten genus is an H_∞ -map. As a corollary, we recover a formula by Dijkgraaf, Moore, Verlinde and Verlinde for the orbifold Witten genus of these symmetric powers. We outline some of the relationship between our power operations and notions from (generalized) Moonshine.

CONTENTS

1. Introduction	2
2. Localization and Fourier expansion	4
2.1. Orbifold loops spaces	5
2.2. Representations of $\mathbb{R}/\mathbb{Z}$	6
2.3. Fourier decomposition of loop bundles	7
3. Thom classes	8
3.1. Devoto's equivariant Tate K-theory	8
3.2. Stringy orbifold Thom classes	9
3.3. The equivariant Witten genus	12
3.4. Integrality	13
3.5. Morita invariance	14
4. Loop spaces of symmetric powers	16
4.1. Actions of wreath products	16
4.2. Loop spaces of symmetric powers	18
4.3. Loop bundles of symmetric products	21
4.4. Iterated symmetric powers	22
5. Power operations	24
5.1. Atiyah power operations	24
5.2. Stringy power operations	27
5.3. Hecke operators	29
5.4. The DMVV-formula, Borchers products and replicability	31
Appendix A. Thom classes and the fixed point formula	34
A.1. Thom classes in equivariant cohomology theories	34
A.2. Atiyah and Segal's Lefschetz fixed point formula	37
References	40

Date: February 8, 2008.

The author was supported by NSF grant DMS-0504539. Most of this paper was written during her stay at MSRI in the spring of 2006.

1. INTRODUCTION

Let M be a compact, closed, complex manifold. Dijkgraaf, Moore, Verlinde and Verlinde proved the following formula about the orbifold elliptic genus of the symmetric powers of M :

$$(1) \quad \sum_{n \geq 0} \phi_{\text{orb}}(M^n // \Sigma_n) t^n = \exp \left(\sum_{m \geq 1} T_m(\phi(M)) t^m \right),$$

where the T_m are Hecke operators. In [Gan06], we proved that such formulas arise whenever the orbifold genus ϕ_{orb} is defined by an H_∞ -map into a cohomology theory with Hopkins-Kuhn-Ravenel theory and (elliptic) power operations. More precisely, let Ell be an elliptic cohomology theory as defined in [AHS01], and let

$$\{\text{Ell}_G \mid G \text{ is a finite group}\}$$

be a compatible family of equivariant versions of Ell . Let $\phi: MU \rightarrow \text{Ell}$ be a map of ring spectra, and assume that we are given equivariant versions $\phi_G: MU_G \rightarrow \text{Ell}_G$ of ϕ . Typically, one can interpret elements $\chi \in \text{Ell}_G(\text{pt})$ as class functions on pairs of commuting elements of G , in which case we say that Ell has a Hopkins-Kuhn-Ravenel character theory. Two notable examples are Borel-equivariant Morava E_2 -theory [HKR00] and Devoto's equivariant Tate K -theory [Dev96].

Definition 1.1. Let $\chi \in \text{Ell}_G(\text{pt})$, where Ell has a Hopkins-Kuhn-Ravenel theory. We define

$$\varepsilon_G(\chi) := \frac{1}{|G|} \sum_{gh=hg} \chi(g, h),$$

and

$$\phi_{\text{orb}} := \varepsilon_G \circ \phi_G.$$

Often, ε_G and thus ϕ_{orb} take values in the coefficient ring Ell_* . Consider the following diagram:

$$(2) \quad \begin{array}{ccc} MU_* & \xrightarrow{\phi} & \text{Ell}_* \\ \oplus P_n \downarrow & & \downarrow \oplus P_n \\ \bigoplus MU_{\Sigma_n} t^n & \xrightarrow{\oplus \phi_{\Sigma_n}} & \bigoplus \text{Ell}_{\Sigma_n} t^n \xrightarrow{\oplus \varepsilon_{\Sigma_n} t^n} \text{Ell}[[t]] \end{array}$$

Here the left vertical arrow is the total power operation in cobordism,

$$P_n([M]) = [M^n \wr \Sigma_n],$$

and t is a dummy variable. The horizontal dotted arrow exists if Ell has a Hopkins-Kuhn-Ravenel theory, and then the composite of the left vertical arrow with the two lower horizontal arrows becomes the left-hand side of the DMVV formula (1).

Assume now that Ell possesses power operations

$$P_n: \text{Ell}_G(X) \rightarrow \text{Ell}_G(X^n).$$

For X the one point space and G the trivial group, this implies that the (right) vertical dotted arrow of the above diagram exists. The map ϕ is called an H_∞ -map if for each n , the equality

$$P_n^{\text{Ell}} \circ \phi = \phi_{\Sigma_n} \circ P_n^{\text{MU}}$$

holds. For such ϕ , the diagram commutes.

If Ell has a Hopkins-Kuhn-Ravenel theory, one can use the P_n to define Hecke operators

$$T_n: \text{Ell}(X) \rightarrow \text{Ell}(X)$$

(cf. [And92]). The P_n are called *elliptic* if this definition of Hecke operators agrees with another one, which is in terms of isogenies [AHS04]. By [Gan06], the composite of the dotted arrows is

$$(3) \quad \sum_{n \geq 0} \varepsilon_{\Sigma_n} (P_n(x)) t^n = \exp \left(\sum_{m \geq 1} T_m(x) t^m \right).$$

If ϕ is an H_∞ -map, this implies formula (1).

In the present paper, we treat the case where Ell is Devoto's equivariant Tate K-theory and ϕ is the equivariant Witten genus. Section 3 introduces the equivariant Witten genus as a map of spectra. Motivated by the work of Lupercio, Uribe and Xicoténcatl on orbifold loop spaces, we define Thom classes in Devoto K-theory for G -equivariant U^2 -bundles, where G is a finite group. We prove that the induced map of spectra realizes the G -equivariant Witten genus defined in [dFLNU06]. We prove that the associated orbifold genus is Morita-invariant and takes values in

$$q^{-d/12} K_{\text{Tate}}(\text{pt}) = q^{-d/12} \mathbb{Z}[[q]],$$

where d is the complex dimension of the manifold.

We proceed to study loop spaces of symmetric powers of orbifolds: If M is a manifold, then a loop in $M^n // \Sigma_n$ is given by n paths in M and a permutation σ , where for each i , the end point of the path γ_i is the starting point of the path $\gamma_{\sigma(i)}$. For each k -cycle of σ , this produces one loop of length k in M . The fact that the loop space of $M^n // \Sigma_n$ is made up from loops in M is the key argument in [DMVV97]. For the symmetric powers of a (global quotient) orbifold $M//G$, a similar picture is true, but the situation is complicated by the additional action of G .

In Section 4, we prove a theorem expressing the orbifold loop spaces of the symmetric powers of $M//G$ in terms of the orbifold loop space of $M//G$ and show that our construction is compatible with iterated symmetric powers. This theorem can also be found in Tamanoi's paper [Tam], which was still unavailable when this paper was written. For trivial G , Lupercio, Uribe, and Xicoténcatl have also given a proof [LUX]. Unlike [DMVV97], their motivation is to study power operations in Adem-Ruan's orbifold cohomology.

In Section 5, we use the analysis of 4 to define equivariant power operations in Devoto K-theory. We compute a formula for the Hecke operators associated to our power operations and conclude that the P_n are elliptic. Interestingly, this agrees exactly with the formula for the twisted Hecke operators in generalized Moonshine found in [Gan]. If G is the trivial group, we recover the operators of [And00].

We proceed to prove that a variant of the equivariant Witten genus is an H_∞ -map. As a corollary, we obtain the Dijkgraaf-Moore-Verlinde-Verlinde formula for our equivariant Witten genus.

The total symmetric power $\text{Sym}_t^{\text{string}}$ is defined as the composite of the two dotted arrows in Diagram (2). It is computed by Formula (3). If $\phi(M)$ is the Witten genus of M , then $\text{Sym}_t^{\text{string}}(\phi(M))$ becomes the right-hand side of the DMVV-formula. At the other hand, we compute $\text{Sym}_t^{\text{string}}(V)$, where V is a complex vector bundle, and find that it equals Witten's

exponential characteristic class

$$\mathrm{Sym}_{\mathbf{t}}^{\mathrm{string}}(\mathbf{V}) = \bigotimes_{k \geq 1} \mathrm{Sym}_{\mathbf{t}^k}(\mathbf{V}),$$

where $\mathrm{Sym}_{\mathbf{t}}$ is the total symmetric power in $\mathbf{K}$ -theory. This sheds some light on the fact that the roles of $\mathbf{t}$ and $\mathbf{q}$ become symmetric, when the right-hand side of (1) is written in its product form

$$\prod_{i,j} \left(\frac{1}{1 - \mathbf{q}^i \mathbf{t}^j} \right)^{c(i,j)}.$$

Here the $c(i)$ are the coefficients of the Witten genus of $\mathbf{M}$,

$$\phi(\mathbf{M}) = \sum c(i) \mathbf{q}^i.$$

Together, the results of this paper fit the DMVV formula for the Witten genus into the picture of equivariant elliptic cohomology explained above.

In [Gan06] we worked in a purely homotopy-theoretic setup and applied our results to the σ -orientation of Morava E_2 -theory, which was shown to be H_∞ in [AHS04]. Morava E -theory has no known geometric definition, and the fact that we could reproduce a DMVV-type formula for the σ -orientation suggests a deep and somewhat mysterious connection between homotopy-theoretic notions such as the $\mathbf{K}(\mathfrak{n})$ -local categories and string-theoretic notions such as orbifold genera.

The results of the paper at hand closely relate to both sides. At the one hand, $\mathbf{K}_{\mathrm{Tate}}$ is an elliptic spectrum, and Devoto's equivariant versions fit into the general framework of equivariant elliptic cohomology and level structures on elliptic curves, axiomatically formalized in [GKV]. Further, when restricted to $\mathrm{MU}\langle 6 \rangle$, the Witten genus becomes the σ -orientation of $\mathbf{K}_{\mathrm{Tate}}$ (cf. [AHS01]).

At the other hand, as we have explained in [AFG], the Witten genus is closely related to the genus considered by Borisov and Libgober. Moreover, Devoto's definition of $\mathbf{K}_{\mathrm{Dev}}$ was inspired by orbifold loop spaces, and so are our definitions of Thom classes and power operations.

1.0.1. Acknowledgements. Many thanks go to Haynes Miller, who generously shared his unpublished work with me. Parts of this paper are closely following his exposition in [Mila] and [Milb]. It is a pleasure to thank Ralph Cohen, Matthew Ando, Haynes Miller, Alex Ghitza, Michael Hopkins and Ernesto Lupercio for long and helpful conversations.

2. LOCALIZATION AND FOURIER EXPANSION

This section introduces some basic definitions. We start with the spaces modelling loops of length k in $\mathbf{M}/\mathbf{G}$. Our definitions are a generalization of the familiar case $k = 1$ (see for example [dFLNU06]). We need to include the case $k > 1$ for technical reasons: it will be necessary when we consider iterated symmetric powers. For the time being the reader is invited to ignore this issue and to set $k = 1$. Throughout the paper we will drop k from the notation if $k = 1$. All our paths are piecewise smooth, and path-spaces carry the compact-open topology.

2.1. Orbifold loops spaces. Let G be a finite group acting smoothly from the right on a manifold M . Let $l = |g|$ be the order of an element g of G , and write $C_g = C_G(g)$ for its centralizer in G .

Definition 2.1. Let k be a natural number. We write C_g^k or $C_G^k(g)$ for the quotient of $C_g \times \mathbb{Z}/kl\mathbb{Z}$ by the normal subgroup generated by (g^{-1}, k) , and $C_g^{\mathbb{R}/k}$ for the quotient of $C_g \times \mathbb{R}/kl\mathbb{Z}$ by the same normal subgroup.

Definition 2.2. For a natural number k , we define

$${}_k\mathcal{P}_g M := \{\gamma: [0, k] \rightarrow M \mid \gamma(k) = \gamma(0)g\}$$

and

$${}_k\mathcal{L}_g M := \text{maps}_{\mathbb{Z}/l\mathbb{Z}}(\mathbb{R}/kl\mathbb{Z}, M)$$

($\mathbb{Z}/l\mathbb{Z}$ -equivariant maps), where $1 \in \mathbb{Z}/l\mathbb{Z}$ acts as k on $\mathbb{R}/kl\mathbb{Z}$ and as g on M .

The centralizer C_g acts on both of these spaces via its action on M , and there is a C_g -equivariant homeomorphism

$$\begin{aligned} {}_k\mathcal{P}_g M &\cong {}_k\mathcal{L}_g M \\ \gamma &\mapsto \gamma * \gamma g * \cdots * \gamma g^{l-1}. \end{aligned}$$




FIGURE 1. A path $\gamma \in \mathcal{P}_g M$ and its image in $\mathcal{L}_g M$. Here $l = 4$.

The group $\mathbb{R}/kl\mathbb{Z}$ (and hence also its subgroup $\mathbb{Z}/kl\mathbb{Z}$) acts on ${}_k\mathcal{L}_g M$ by rotation of the loops. Note that the actions of $k \in \mathbb{R}/kl\mathbb{Z}$ and $g \in C_g$ agree. Hence we have actions of $C_g^{\mathbb{R}/k}$ and C_g^k on ${}_k\mathcal{L}_g M$. Using the above homeomorphism, we obtain $C_g^{\mathbb{R}/k}$ and C_g^k -actions on ${}_k\mathcal{P}_g M$. Throughout the paper we will use this homeomorphism to identify ${}_k\mathcal{L}_g M$ with ${}_k\mathcal{P}_g M$ and, by abuse of notation, we will write ${}_k\mathcal{L}_g M$ for both.

Definition 2.3. Let ${}_k\mathcal{L}(M//G)$ be the disjoint union over the conjugacy classes of G

$${}_k\mathcal{L}(M//G) := \coprod_{[g]} {}_k\mathcal{L}_g M // C_g^k.$$

We will often think of ${}_k\mathcal{L}(M//G)$ as the groupoid of k open strings in $M//G$ joining together to form one long closed string, where the order of the k strings does not matter (see Figure 2). We are now ready to define loop vector bundles over orbifold loop spaces.

Definition 2.4. Let $\mathcal{G}$ be one of the groups $U(n)$ or $Spin(2n)$. An (orbifold) $\mathbb{S}^1$ -equivariant $\mathcal{L}\mathcal{G}$ -vector bundle over ${}_k\mathcal{L}_g M // C_g^k$ is a (typically infinite dimensional) vector bundle $\mathcal{V}$ over ${}_k\mathcal{L}_g M$ with structure group ${}_k\mathcal{L}\mathcal{G}$, which is $C_g^{\mathbb{R}/k}$ -equivariant as ${}_k\mathcal{L}\mathcal{G}$ -bundle (i.e., the groups




FIGURE 2. An element $\gamma = \gamma_1 * \cdots * \gamma_5 \in {}_5\mathcal{LM}$ for trivial G .

act by ${}_{\text{kl}}\mathcal{LG}$ -bundle automorphisms), such that the action of $\mathbb{R}/\text{kl}\mathbb{Z}$ on $\mathcal{V}$ intertwines with its action on ${}_{\text{kl}}\mathcal{LG}$.

Example 2.5. Let $V \in \text{Vect}(\underline{X})$ be a $\mathcal{G}$ -vector bundle on the orbifold $\underline{X}$. View V as an orbifold over $\underline{X}$. Then the loop orbifold $\mathcal{LV}$ of V is an orbifold $\mathbb{S}^1$ -equivariant $\mathcal{LG}$ -vector bundle over $\mathcal{LX}$.

Definition 2.6. Two ${}_{\text{l}}\mathcal{LG}$ -vector bundles $\mathcal{V}$ and $\mathcal{W}$ over a space X are called *densely isomorphic* if there are dense sub-bundles $\mathcal{V}' \subseteq \mathcal{V}$ and $\mathcal{W}' \subseteq \mathcal{W}$ and an isomorphism of vector bundles with structure group $\mathcal{LG}$ between $\mathcal{V}'$ and $\mathcal{W}'$. If in addition $\mathcal{V}$ and $\mathcal{W}$ are G -equivariant bundles over a G -space X , then $\mathcal{V}$ is densely isomorphic to $\mathcal{W}$ as a loop bundle with this extra structure, if the inclusions and the vector bundle isomorphisms preserve this extra structure. We will denote dense isomorphisms by “ $\cong$ ” and often drop the word “*densely*”.

2.2. Representations of $\mathbb{R}/l\mathbb{Z}$. Similarly to the non-equivariant case, there is a Fourier decomposition principle for loop bundles that are restricted to the constant loops. In the orbifold case, fractional powers of $\mathfrak{q}$ can occur in these decompositions. The reason for this phenomenon is that in the Fourier series loops of length $|g|$ need to be rescaled with a factor $1/|g|$.

Definition 2.7. Let $\mathfrak{q}^{\frac{1}{l}}$ denote the complex representation of $\mathbb{R}/l\mathbb{Z}$ obtained by precomposing the fundamental representation of $\mathbb{S}^1$ with the map

$$\begin{aligned} \mathbb{R}/l\mathbb{Z} &\rightarrow \mathbb{R}/\mathbb{Z} \\ t &\mapsto t/l. \end{aligned}$$

Note that

$$1 \in \mathbb{Z}/l\mathbb{Z} \subseteq \mathbb{R}/l\mathbb{Z}$$

acts on $\mathfrak{q}^{\frac{1}{l}}$ by multiplication with a primitive l^{th} root of unity, ζ_l . Recall that all non-trivial irreducible real representations of $\mathbb{R}/l\mathbb{Z}$ are of the form

$$\mathfrak{q}_{\mathbb{R}}^{\frac{j}{l}} := \text{res}_{|\mathbb{R}}^{\mathbb{C}} \mathfrak{q}^{\frac{j}{l}},$$

where $\text{res}_{|\mathbb{R}}^{\mathbb{C}} \rho$ denotes the underlying real vector space of ρ . Further, $\mathfrak{q}^{\frac{i}{l}} \cong_{\mathbb{R}} \mathfrak{q}^{\frac{j}{l}}$ if and only if $i = \pm j$. Let

$$\rho: \mathbb{R}/l\mathbb{Z} \rightarrow \mathcal{G} \rightarrow \text{SO}(2n)$$

be a $\mathcal{G}$ -representation of $\mathbb{R}/\mathbb{L}\mathbb{Z}$ with underlying vector space V . Then V decomposes into a direct sum of complex/Spin representations

$$(4) \quad V \cong V_0 \oplus \bigoplus_{j \geq 1} V_j q^{\frac{j}{4}},$$

where $V_j q^{\frac{j}{4}}$ denotes the summand with rotation number j . This notation is motivated as follows: For $j \geq 1$, we can endow the underlying vector space V_j of $V_j q^{\frac{j}{4}}$ with a complex structure, where multiplication with i is defined as the action of $\frac{1}{4j}$. Then

$$V_j q^{\frac{j}{4}} = \text{res}_{\mathbb{R}}^{\mathbb{C}} \left(V_j \otimes_{\mathbb{C}} \left(q^{\frac{1}{4}} \right)^{\otimes j} \right).$$

Warning: if $\mathcal{G}$ is $U(n)$, then V decomposes as a complex representation

$$V \cong_{\mathbb{C}} \bigoplus_{j \in \mathbb{Z}} W_j q^{\frac{j}{4}},$$

but (4) endowed with the complex structure defined by the rotations is

$$(5) \quad V \cong_{\mathbb{R}} W_0 \oplus \bigoplus_{j \geq 1} (\overline{W}_{-j} \oplus W_j) q^j$$

2.3. Fourier decomposition of loop bundles. Let $\mathcal{V}$ be an orbifold $\mathbb{S}^1$ -equivariant $\mathcal{LG}$ vector bundle over ${}_{\kappa}\mathcal{L}_g M // C_g^k$. The condition that the action of $\mathbb{R}/k\mathbb{L}\mathbb{Z}$ on $\mathcal{V}$ intertwines with that of ${}_{\kappa}\mathcal{L}\mathcal{G}$ spells out to the following: they combine to a (left-) action of $\mathbb{R}/k\mathbb{L}\mathbb{Z} \ltimes_{\kappa} \mathcal{LG}$ on $\mathcal{V}$, where

$$(t, \gamma') \in \mathbb{R}/k\mathbb{L}\mathbb{Z} \ltimes_{\kappa} \mathcal{LG}$$

maps $v \in \mathcal{V}_{\gamma}$ to

$$(t, \gamma') \cdot v = t \cdot (\gamma' \cdot v) = (t \cdot \gamma')(t \cdot v) \in \mathcal{V}_{t \cdot \gamma}.$$

Restricting to constant loops, we get an action of $C_g^{\mathbb{R}/k} \times \mathcal{G}$ on $\mathcal{V}|_{M^g}$, where $\mathbb{R}/k\mathbb{L}\mathbb{Z} \times \mathcal{G}$ acts fiber preserving. Therefore, $\mathcal{V}|_{M^g}$ decomposes into a direct sum of $\mathbb{R}/k\mathbb{L}\mathbb{Z}$ -representations

$$\mathcal{V}|_{M^g} \cong_{\mathbb{R}} V_0 \oplus \bigoplus_{j \geq 1} V_j q^{\frac{j}{k\mathbb{L}}}$$

as above, where $V_j q^{\frac{j}{k\mathbb{L}}}$ is a (finite dimensional) $C_g^{\mathbb{R}/k}$ -equivariant $\mathcal{G}$ -vector bundle over M^g . The actions of $g \in C_g$ and $k \in \mathbb{R}/k\mathbb{L}\mathbb{Z}$ on V_j agree, so that g acts on V_j as (complex) multiplication by $\zeta_{\mathbb{L}}^j$.

Example 2.8. Let V be a finite dimensional real vector space, and recall (e.g. from [HBJ92, 6]) that there is a dense isomorphism of real $\mathbb{R}/\mathbb{L}\mathbb{Z}$ -representations

$$\text{maps}(\mathbb{R}/\mathbb{L}\mathbb{Z}, V) \cong V \oplus \bigoplus_{j \geq 1} V_{\mathbb{C}} q^{\frac{j}{4}},$$

given by the Fourier expansion principle. Here $V_{\mathbb{C}}$ denotes the complexification of V . Let now $\langle g \rangle$ act on V , where $|g| = \mathbb{L}$. Note that $1 \in \mathbb{R}/\mathbb{L}\mathbb{Z}$ acts by multiplication with $\zeta_{\mathbb{L}}^j$ on the j^{th} summand of the right-hand side. A loop is $\mathbb{Z}/\mathbb{L}\mathbb{Z}$ -equivariant, if and only if its Fourier expansion $\sum v_j q^j$ satisfies

$$(\forall j \geq 0) \quad v_j g = \zeta_{\mathbb{L}}^j v_j.$$

Hence we have

$$\mathrm{maps}_{\mathbb{Z}/l}(\mathbb{R}/l\mathbb{Z}, V) \cong V_1 \oplus \bigoplus_{j \geq 1} (V_{\mathbb{C}})_{\zeta_l^j} q^{\frac{j}{l}}.$$

Here $(V_{\mathbb{C}})_{\zeta}$ denotes the ζ -eigenspace of the complexification of V , and similarly, V_1 denotes the vectors of V fixed by g .

Example 2.9. Let $\mathcal{V} = \mathcal{L}(V//G)$ be the orbifold loop bundle of a G -equivariant $\mathcal{G}$ -vector bundle over M . Let l be the order of $g \in G$, and let m be a point in M^g . We have

$$\mathcal{L}_g V|_m \cong \mathrm{maps}_{\mathbb{Z}/l}(\mathbb{R}/l\mathbb{Z}, V|_m)$$

and hence

$$\mathcal{L}_g V|_{M^g} \cong V_1 \oplus \bigoplus_{j \geq 1} (V_{\mathbb{C}})_{\zeta_l^j} q^{\frac{j}{l}},$$

where $(V_{\mathbb{C}})_{\zeta}$ denotes the ζ -eigenbundle of the action of g on $V_{\mathbb{C}}|_{M^g}$. (Compare also [dFLNU06, (5.1.1)].)

3. THOM CLASSES

3.1. Devoto's equivariant Tate K-theory. Taking g -fixed points is a functor

$$(-)^g: G\text{-spaces} \rightarrow C_g\text{-spaces}$$

which preserves cofibre sequences, wedges and weak equivalences. Therefore

$$(6) \quad X \mapsto \bigoplus_{[g]} K_{C_g}(X^g) \llbracket q^{\frac{1}{|g|}} \rrbracket$$

satisfies the axioms of a G -equivariant cohomology theory in [May96]. If g is conjugate to h in G (denoted $g \sim h$) there is a canonical isomorphism

$$K_{C_g}(X^g) \cong K_{C_h}(X^h).$$

Thus (6) is, up to canonical isomorphism, independent of choices of the representatives.

Definition 3.1. We define the Devoto equivariant Tate K-theory rings (for strings of length r) to be the subring of

$$K_{\mathrm{Dev}, G, r}(X) := \bigoplus_{[g]} K_{C_g^r}(X^g) \llbracket q^{\frac{1}{r|g|}} \rrbracket$$

with the property that $1 \in \mathbb{Z}/r|g|\mathbb{Z} \subseteq C_g^r$ acts by complex multiplication with $\zeta_{r|g|}^j$ on the coefficient of $q^{\frac{j}{r|g|}}$. If $r = 1$, we will drop it from the notation.

Remark 3.2. For $r = 1$, the condition of the definition should be compared to Condition (a) of the generalized Moonshine axioms: the Möbius transformation $\tau \mapsto \tau + 1$ (sending $q^{\frac{j}{|g|}}$ to $q^{\frac{j}{|g|}} \zeta_{|g|}^j$) should correspond to the transition from (g, h) to (g, gh) , i.e., to the action of g on the summand corresponding to $[g]$.

Definition 3.3. Let $\alpha: H \rightarrow G$ be a map of finite groups, and let $x \in K_{\text{Dev}}(X//G)$. Then

$$(\text{res}|_{\alpha} x)_{[h]} = x_{[\alpha(h)]}.$$

Let $x \in K_{\text{Dev},G}(X)$ and $y \in K_{\text{Dev},H}(Y)$. Then the external tensor product of x and y is the element

$$x \boxtimes y \in K_{\text{Dev},G \times H}(X \times Y)$$

defined by

$$(x \boxtimes y)_{[g,h]} := x_{[g]} \boxtimes y_{[h]} \in K_{C_G(g) \times C_H(h)}(X^g \times Y^h) \llbracket q^{\frac{1}{N}} \rrbracket,$$

where N is the smallest common multiple of $|g|$ and $|h|$, $\boxtimes$ is the external tensor product of vector bundles, and $q^r \boxtimes q^s := q^{r+s}$. If $G = H$, the G -equivariant external tensor product is defined as

$$x \boxtimes y := \text{res}|_{\delta}(x \boxtimes y),$$

where δ denotes the diagonal inclusion of G in $G \times G$. If in addition $X = Y$, then the internal tensor product of x and y is defined to be the pullback along the diagonal d of $X \times X$ of $x \boxtimes y$:

$$x \otimes y := d^*(x \boxtimes y).$$

3.2. Stringy orbifold Thom classes. Let G be a finite group, let g be an element of G , and let l be the order of g . Let V be a G -equivariant $\text{Spin}(2d)$ -vector bundle over a G -space X . By Example 2.9, we have

$$(7) \quad \mathcal{L}_g V|_{X^g} \cong V_1 \oplus \bigoplus_{j \geq 1} (V_{\mathbb{C}})_{\zeta_l^j} q^{\frac{j}{l}}.$$

Assume that X^g is connected. The bundle V_1 is the bundle of g -fixed points of $V|_{X^g}$, and the g -fixed points of the Thom space of V are, as a C_g -space, homeomorphic to the Thom space $(X^g)^{V_1}$ of V_1 .

Assume first that g acts trivially on $V|_{X^g}$, then $V_1 = V|_{X^g}$, and (7) contains only the terms with integral powers of q . This case was treated in [Milb], we shortly recall the discussion there: Since V is oriented, $V_{\mathbb{C}}$ is an SU -vector bundle. Hence it has a $\text{Spin}(4d)$ structure as well as a complex structure with trivial determinant bundle

$$\det V_{\mathbb{C}} = 1 \in K_{C_g}(X^g),$$

and we have

$$e_{C_g}^{\hat{\Lambda}}(V_{\mathbb{C}}|_{X^g}) = e_{C_g}^{\text{Td}}(V_{\mathbb{C}}|_{X^g}).$$

Formula (20) applied to the $\mathbb{R}/\mathbb{U}(1)$ -action suggests to define the $C_g \times \mathbb{R}/\mathbb{U}(1)$ -equivariant Thom class of $\mathcal{L}_g V|_{X^g}$ as

$$u_{C_g}^{\hat{\Lambda}}(V) \cdot \prod_{j \geq 1} q^{-dj} \cdot \wedge_{-q^j} V_{\mathbb{C}}.$$

Here the exterior powers are exterior powers of C_g -representations. It is customary [Seg88] to use a ζ -function renormalization to replace the divergent sum $\sum_{j > 0} j$ in the exponent of q with $\zeta(-1) = -1/12$, and to define

$$u_g^{\text{string}}(V) := q^{\frac{d}{12}} \cdot u_{C_g}^{\hat{\Lambda}}(V) \cdot \prod_{j=1}^{\infty} \wedge_{-q^j} V_{\mathbb{C}}.$$

Next, assume that $\mathfrak{g}$ acts fixed point free on $V|_{X^g}$ and that -1 is not an eigenvalue of $\mathfrak{g}$. For $j = 1, \dots, \lfloor \frac{l}{2} \rfloor$, let $\theta_j = 2\pi \frac{j}{l}$, and let V_j be the subspace of V on which $\mathfrak{g}$ acts by rotations with angle θ_j . We equip V with the unique complex structure such that $\mathfrak{g}$ acts on V_j by multiplication with $e^{i\theta_j}$. Then $V_j \otimes \mathbb{C}$ is of the form

$$V_j \otimes \mathbb{C} \cong V_j \oplus \overline{V_j}.$$

Setting $V_{-j} := \overline{V_j}$, we have

$$\mathcal{L}V|_{X^g} \cong \bigoplus_{j=1}^{\lfloor \frac{l}{2} \rfloor} V_j q^{\frac{j}{l}} \oplus \bigoplus_{k=1}^{\infty} \bigoplus_{j=-\lfloor \frac{l}{2} \rfloor}^{\lfloor \frac{l}{2} \rfloor} V_j q^{k+\frac{j}{l}}.$$

For fixed k ,

$$\begin{aligned} \det \left(\bigoplus_{j=-\lfloor \frac{l}{2} \rfloor}^{\lfloor \frac{l}{2} \rfloor} V_j q^{k+\frac{j}{l}} \right) &\cong \bigotimes_{j=-\lfloor \frac{l}{2} \rfloor}^{\lfloor \frac{l}{2} \rfloor} q^{(k+\frac{j}{l})d_j} \cdot \det(V_j) \\ &\cong q^{2dk} \cdot \det(V_{\mathbb{C}}). \end{aligned}$$

Here $d_j = \dim_{\mathbb{C}} V_j$, and

$$\sum_{j=-\lfloor \frac{l}{2} \rfloor}^{\lfloor \frac{l}{2} \rfloor} d_j(k + \frac{j}{l}) = 2k \left(\sum_{j=1}^{\lfloor \frac{l}{2} \rfloor} d_j \right) = 2dk,$$

while $\det V_{\mathbb{C}}$ is trivial. In a similar manner, the determinant line bundle of the first $\lfloor \frac{l}{2} \rfloor$ summands becomes

$$q^{\frac{\text{age}(\mathfrak{g})}{2}} \cdot \det(V)$$

with

$$\text{age}(\mathfrak{g}) = 2 \cdot \sum_{j=1}^{\lfloor \frac{l}{2} \rfloor} \frac{j}{l} d_j.$$

Note that $\det(V)$ is not trivial, but has a square root.

As above, Formula (20) suggests to define the $C_g \times \mathbb{R}/\mathbb{Z}$ -equivariant Euler/Thom class of $\mathcal{L}_g V|_{X^g}$ as

$$q^{\frac{d}{12} - \frac{\text{age}(\mathfrak{g})}{2}} \left(\sqrt{\det V} \right)^{-1} \cdot \prod_{j=1}^{\infty} \wedge_{-q^{\frac{j}{l}}} \left((V_{\mathbb{C}})_{\zeta_l^j} \right).$$

If we want to allow the eigenvalue -1 , we need to assume that the eigenspace $V_{\frac{l}{2}}$ of the action of -1 on V has a complex structure. Then we can make sense of the expression $(\sqrt{\det V})$, and the above discussion goes through.

This is for example the case if V is a G -equivariant U^2 -bundle. Note that in that case we have two complex structures on the other V_j – the given complex structure and the auxiliary one we defined above. We choose to work with the latter. This is an issue similar to the one in the warning after Formula (4). Write V_1 for the subspace of $V|_{X^g}$ that is fixed by the action of $\mathfrak{g}$, write N^g for its orthogonal complement, and note that

$$u_{C_g}^{\hat{A}}(V_1)(\sqrt{\det N^g})^{-1} = u_{C_g}^{\text{Td}}(V_1)(\sqrt{\det V_1})^{-1}(\sqrt{\det N^g})^{-1} = u_{C_g}^{\text{Td}}(V_1)(\sqrt{\det V})^{-1}.$$

Together with the above discussion, this motivates the following definition:

Definition 3.4. Let V be a G -equivariant $U^2(d)$ -vector bundle over a compact G -space X . Let l be the order of the element $g \in G$. Assume that X^g is connected. We define

$$u_g^{\text{string}}(V) := q^{\frac{d}{12} - \frac{\text{age}(g)}{2}} \cdot u_{C_g}^{\text{Td}}(V_1) \cdot (\sqrt{\det V})^{-1} \cdot \prod_{j=1}^{\infty} \Lambda_{-q^j} \left((V_{\mathbb{C}})_{\zeta_l^j} \right)$$

and set

$$u^{\text{string}}(V) := \sum_{[g]} u_g^{\text{string}}(V).$$

This is an element of $K_{\text{Dev}, G}(X^V)[q^{\pm \frac{1}{12}}]$. If X^g is not connected, one sums over its connected components.

Proposition 3.5. *The $u_G^{\text{string}}(V)$ satisfy the axioms of Definition A.1.*

PROOF : The naturality axiom and the change of groups axiom are straight-forward. Multiplicativity follows from that of u^{Td} and from the multiplicative properties of $\sqrt{\det(-)}$ and of the total exterior power Λ_t . Further, Λ_{-t} is invertible with inverse $\text{Sym}_t(V)$, which implies the periodicity axiom. $\square$

3.2.1. *Comparison to the work of Kitchloo and Morava.* Let G be the trivial group, and let V be a complex vector bundle. Using (5) and the fact that

$$\begin{aligned} \Lambda_{-1}(Vq^{-k} \oplus Vq^k) &\cong \det(-Vq^{-k}) \cdot \Lambda_{-1}(Vq^k) \\ &\cong q^{dk}(\det V)^{-1} \cdot \Lambda_{-1}(Vq^k), \end{aligned}$$

we see that if $\det(V)$ is trivial, a similar discussion as above suggests the definition

$$u_{\text{Td}}^{\text{string}}(V) := q^{-\frac{1}{12}} u_{\text{Td}}(V) \cdot \prod_{k \geq 1} \Lambda_{-q^k} V_{\mathbb{C}}.$$

The periodicity element $\alpha = u_{\text{Td}}^{\text{string}}(\mathbb{C})$ (c.f. (14)) is

$$\alpha = \left(q^{-\frac{1}{12}} \cdot \prod_{j \geq 1} (1 - q^j)^2 \right) \cdot \beta,$$

where β is the Bott-element. Hence the renormalized Thom class in degree $2d$

$$\begin{aligned} \tau_{\text{Td}}^{\text{string}}(V) &= u_{\text{Td}}^{\text{string}}(V) \cdot \alpha^{-d} \\ &= \tau^{\text{Td}}(V) \cdot \prod_{j \geq 1} \frac{\Lambda_{-q^j} V_{\mathbb{C}}}{(1 - q^j)^{2d}} \end{aligned}$$

is the Thom class defined in [AM01], [KM, 4.1].

3.3. The equivariant Witten genus.

Corollary 3.6 (of Proposition 3.5). *There is a unique map of G -spectra*

$$\phi_G: MU_G^2 \rightarrow K_{\text{Dev}, G}$$

taking Thom classes to Thom classes. The induced genus

$$\hat{A}_G^{\text{string}}(M) := \phi_G(M)$$

of a G -equivariant U^2 manifold M of dimension $2n$ is given by the formula

$$\left(\hat{A}_G^{\text{string}}(M) \right)_{[g]} = q^{-\frac{d}{12} + \frac{\text{age}(g)}{2}} \cdot \text{Td}_{C_g} \left(M^g; \sqrt{TM} \cdot \prod_{j \geq 1} \text{Sym}_{q^j} \left((TM^C)_{\zeta_l^j} \right) \right),$$

where l is the order of g , and we sum over the connected components of M^g .

PROOF : The existence and uniqueness of ϕ_G follow from Proposition A.4. Further, assuming M^g connected,

$$\begin{aligned} \left(\hat{A}_G^{\text{string}}(M) \right)_{[g]} &\cong \left(\pi_l^{\text{string}}(1) \right)_{[g]} \\ &\cong \pi_{C_g!}^{\text{Td}} \left(\left(q^{\frac{d}{12} - \frac{\text{age}(g)}{2}} (\sqrt{TM})^{-1} \prod_{j \geq 1} \wedge_{-q^j} (TM^C)_{\zeta_l^j} \right)^{-1} \right) \\ &\cong q^{-\frac{d}{12} + \frac{\text{age}(g)}{2}} \cdot \text{Td}_{C_g} \left(M^g, \sqrt{TM} \prod_{j \geq 1} \text{Sym}_{q^j} (TM^C)_{\zeta_l^j} \right), \end{aligned}$$

where the first and last equation are (14), and the second equation follows from Definition 3.4. $\square$

It is traditional to express equivariant genera in the following more explicit and somewhat more complicated form.

Corollary 3.7. *Let $h \in C_g$, and $k = |h|$. Assume M^g to be connected. Then*

$$\begin{aligned} \hat{A}_G^{\text{string}}(M)_{[g]}(h) &= \pm q^{-\frac{d}{12} + \frac{\text{age}(g)}{2}} e^{i\pi \text{age}(h)} \int_{M^{g,h}} e^H(M^{g,h}) \cdot \left(\prod_{\tilde{x}_i} e^{\frac{\tilde{x}_i}{2}} \right) \\ &\quad \cdot \left[\prod_{s, x_i} (1 - \zeta_k^s e^{x_i}) \prod_{j \geq 1} \prod_{s, y_i} (1 - \zeta_k^s e^{y_i} q^j) \right]^{-1}, \end{aligned}$$

where e^H denotes the Euler class in ordinary cohomology, s runs from 0 to $k-1$, the $\tilde{x}_i$ run over the Chern roots of TM , the x_i are the Chern roots of $TM_{g=1, h=\zeta_k^s}$, and the y_i are the Chern roots of the simultaneous eigenspaces $(TM_C)_{g=\zeta_l^j, h=\zeta_k^s}$. (All bundles are over the double fixed points $M^{g,h}$.)

PROOF : By (17), we have

$$(8) \quad \text{Td}_{C_g}(M^g; a)(h) = \int_{M^{g,h}} e^H(M^{g,h}) \cdot \text{ch}(a(h)) \cdot \prod_{s, x_i} \frac{1}{1 - \zeta_k^s e^{x_i}},$$

where s runs from 0 to $k-1$, and the x_i are the Chern roots of $\mathbf{TM}_{g=1, h=\zeta_k^s}$. We have

$$\mathrm{ch}(\sqrt{\mathbf{TM}}(h)) = \pm \prod_{s, \tilde{x}_i} \zeta_{2k}^s e^{\frac{\tilde{x}_i}{2}} = \pm e^{i\pi \mathrm{age}(h)} \cdot \prod_{\tilde{x}_i} e^{\frac{\tilde{x}_i}{2}},$$

where the $\tilde{x}_i$ are the Chern roots of $(\mathbf{TM})_{h=\zeta_k^i}$. The Chern character of the remaining factor, evaluated at h , computes to

$$\prod_{j \geq 1} \prod_{s=0}^{k-1} \prod_{y_i} \left(1 - \zeta_k^s e^{y_i} q^{\frac{j}{l}}\right),$$

where the y_i are the Chern roots of the simultaneous eigenspaces $(\mathbf{TM}_{\mathbb{C}})_{g=\zeta_l^i, h=\zeta_k^s}$. $\square$

Corollary 3.7 should be compared to the calculations on the last page of [dFLNU06]. Basically, our discussion starts, once they replace their variable $\hat{q}$ with its monodromy $2\pi i$. For our purposes, the double loop space picture is not necessary. It will, however, become very important when one wants to study modularity properties, i.e., the “moonshine-like” behavior discussed in [Dev96]: We already observed that replacing the pair of commuting elements (g, h) with (g, gh) corresponds to replacing τ with $\tau + 1$, where $q = e^{2\pi i \tau}$. This is half of the first Moonshine condition, the other half demands that switching to (h^{-1}, g) should correspond to $\tau \mapsto -\frac{1}{\tau}$. In the picture described in the beautiful paper [dFLNU06], the elements g and h describe the monodromy of a G -principal bundle P over a torus, where P maps equivariantly to M . As described in [FQ93], [Gan], the transformation behaviour of generalized Moonshine corresponds to a different choice of circles, generating the same torus. Our approach sees the first circle, but not the second one. This is the reason why we can explain the first transformation geometrically, but not the second one.

Definition 3.8. Let M be a G -equivariant $U^2(d)$ -manifold. Then the orbifold Witten genus of $M//G$ is defined by

$$\hat{A}_{\mathrm{orb}}^{\mathrm{string}}(M//G) := \frac{1}{|G|} \sum_{gh=hg} \left(\hat{A}_G^{\mathrm{string}}(M)_{[g]} \right) (h),$$

where the sum is over all pairs of commuting elements of G .

One of the advantages of working with our setup is that the proofs of integrality and Morita invariance become fairly straightforward.

3.4. Integrality.

Proposition 3.9. *The orbifold genus $\hat{A}_{\mathrm{orb}}^{\mathrm{string}}$ takes values in $\mathbb{Z}[[q]][q^{\pm \frac{1}{12}}]$.*

PROOF : Let $g \in G$ have order l . Then $\hat{A}_G^{\mathrm{string}}(M)_{[g]}$ is an element of $R(C_g)[[q^{\frac{1}{l}}]][q^{\pm \frac{1}{12}}]$, and

$$\begin{aligned} \hat{A}_{\mathrm{orb}}^{\mathrm{string}}(M//G) &= \sum_{[g]} \frac{1}{|C_g|} \sum_{h \in C_g} \hat{A}_G^{\mathrm{string}}(M)_{[g]}(h) \\ &= \sum_{[g]} \langle \hat{A}_G^{\mathrm{string}}(M)_{[g]}, 1 \rangle_{C_g}. \end{aligned}$$

Here $\langle -, - \rangle_{C_g}$ denotes the inner product in $R(C_g)$, so that

$$\begin{aligned} \left\langle \sum V_j q^{\frac{j}{t}}, 1 \right\rangle_{C_g} &:= \sum \langle V_j, 1 \rangle_{C_g} q^{\frac{j}{t}} \\ &= \sum \dim_{\mathbb{C}} \left(V_j^{C_g} \right) q^{\frac{j}{t}} \end{aligned}$$

computes the dimensions of the trivial summands of the V_j . Fix g for the remainder of the proof, and assume, for simplicity, that M^g is connected. The element $g \in C_g$ acts by multiplication with ζ_l^k on the coefficient of $q^{\frac{k}{t}}$ in

$$\prod_{j \geq 1} \Lambda_{-q^{\frac{j}{t}}}(\mathcal{TM}^{\mathbb{C}})_{\zeta_l^j}.$$

Further, g acts by multiplication with $\pm e^{-i\pi \text{age}(g)}$ on

$$q^{-\frac{\text{age}(g)}{2}} u_{\text{Td}}(\mathcal{TM}^g) \cdot (\sqrt{\det \mathcal{TM}})^{-1}.$$

To determine the sign, note that g acts trivially on $u_{\text{Td}}(M^g)$ and by multiplication with $e^{2i\pi \text{age}(g)}$ on $\det \mathcal{TM}$, and that the sign depends on the choice of an identification of $\text{Spin}(2)$ with $\mathbb{U}(1)$. For our calculation of $\hat{A}^{\text{string}}(M)_{[g]}$, we fix this identification in such a way that the sign becomes positive. Then only the coefficients of integral powers of q in

$$\pi_{\text{Td},!}^{M^g} \left(\left(q^{-\frac{\text{age}(g)}{2}} \cdot \sqrt{\det \mathcal{TM}}^{-1} \cdot \prod_{j \geq 1} \Lambda_{-q^{\frac{j}{t}}}(\mathcal{TM}^{\mathbb{C}})_{\zeta_l^j} \right)^{-1} \right)$$

contribute trivial summands. If we had chosen to work with the other choice of identification, we would have to multiply the above result for $\hat{A}^{\text{string}}(M)_{[g]}$ with -1 , which would not destroy the integrality. $\square$

3.5. Morita invariance. For detailed definitions of the notions of *inertia groupoid* and *orbifold vector bundle* used in this section we refer the reader to [Moe02]. Let $\mathbf{G}$ be an orbifold groupoid, and let $\Lambda(\mathbf{G})$ be its inertia groupoid. Recall that

$$\Lambda(\mathbf{G})_0 = \text{eq}(s, t) \quad \text{and} \quad \Lambda(\mathbf{G})_1 = \text{eq}(s, t) \times_{G_0} G_1$$

(here $\text{eq}(s, t)$ stands for the equalizer of the source and target of $\mathbf{G}$), and consider the section

$$\begin{aligned} \mathbf{A}: \text{eq}(s, t) &\longrightarrow \text{eq}(s, t) \times_{G_0} G_1 \\ g &\longmapsto (g, g). \end{aligned}$$

One checks that $\mathbf{A}$ is a natural transformation from the identity map of $\Lambda(\mathbf{G})$ to itself. Let V be a vector bundle over the groupoid $\Lambda(\mathbf{G})$. By definition, V is a vector bundle over $\Lambda(\mathbf{G})_0$ together with an isomorphism

$$\mu: t^*V \rightarrow s^*V$$

over $\Lambda(\mathbf{G})_1$ such that a few diagrams commute. The section $\mathbf{A}$ pulls back μ to a (fiber preserving) automorphism of

$$V = \mathbf{A}^*t^*V = \mathbf{A}^*s^*V$$

over $\Lambda(\mathbf{G})_0$, which turns out to be an automorphism of V as orbifold vector bundle. All of the above is preserved by equivalences of orbifold groupoids $\mathbf{G} \simeq \mathbf{H}$.

Example 3.10. Let $\mathbf{G} = \mathbf{M} \rtimes \mathbf{G}$ be the translation groupoid of the action of a finite group $\mathbf{G}$ on a manifold $\mathbf{M}$, and let $\mathbf{V}$ be a $\mathbf{G}$ -equivariant vector bundle on $\mathbf{M}$. Then $\Lambda(\mathbf{G})$ is equivalent to the disjoint union over the conjugacy classes of $\mathbf{G}$ of the translation groupoids of the actions of the centralizers on the fixed point loci:

$$\Lambda(\mathbf{G}) \simeq \coprod_{[g]} \mathbf{M}^g \rtimes \mathbf{C}_g.$$

Under this equivalence, the map $\mathbf{A}$ corresponds to the coproduct of the maps

$$\begin{aligned} \mathbf{M}^g &\rightarrow \mathbf{M}^g \times \mathbf{C}_g \\ \mathbf{m} &\mapsto (\mathbf{m}, g), \end{aligned}$$

and $\mathbf{A}^*(\mu)|_{\mathbf{M}^g}$ becomes multiplication with g on $\mathbf{V}|_{\mathbf{M}^g}$.

Let now $\underline{\mathbf{X}}$ be a compact complex orbifold and $\mathbf{TX}$ its tangent bundle. After pulling back along the “inclusion of the fixed point locus” map

$$i: \Lambda \underline{\mathbf{X}} \rightarrow \underline{\mathbf{X}},$$

$\mathbf{TX}$ decomposes into eigenbundles of $\mathbf{A}^*\mu$,

$$\mathbf{TX}|_{\Lambda \underline{\mathbf{X}}} \cong \bigoplus_{\zeta} (\mathbf{TX})_{\zeta},$$

where the sum is over the eigenvalues of $\mathbf{A}^*\mu$. It follows from Example 3.10 and the analogous statement for the equivariant case that locally (and hence globally) we have $(\mathbf{TX})_1 \cong \mathbf{T}\Lambda \underline{\mathbf{X}}$. Let $\mathbf{L}$ be the smallest common multiple of the orders of all the elements of the stabilizers of $\underline{\mathbf{X}}$, and recall that the connected components of an orbifold are well-defined (they correspond to the connected components of its quotient space) and that it is still true for orbifolds that the dimension of a vector bundle is constant over each connected component. Hence, over each connected component of $\underline{\mathbf{X}}$,

$$q^{-\frac{d}{12} + \frac{\text{age}(\mathbf{A}^*(\mu))}{2}} \prod_{j \geq 1} \text{Sym}_{q^{\frac{j}{\mathbf{L}}}} (\mathbf{TX}^{\mathbb{C}})_{\zeta_{\mathbf{L}}^j}$$

is a well defined element of $\mathbf{K}_{\text{orb}}(\Lambda \underline{\mathbf{X}})[[q^{\pm \frac{1}{12}}]]$. Here $\mathbf{TX}^{\mathbb{C}}$ is restricted to $(\Lambda \underline{\mathbf{X}})$.

Let $\mathcal{A}$ be an abelian sheaf over the orbifold $\underline{\mathbf{X}}$. Recall that the orbifold cohomology of $\underline{\mathbf{X}}$ with coefficients in $\mathcal{A}$, denoted

$$\mathbf{H}^*(\underline{\mathbf{X}}, \mathcal{A}),$$

is defined as the cohomology of the complex $\Gamma_{\text{inv}}(\mathcal{I}^{\bullet})$, where

$$0 \rightarrow \mathcal{A} \rightarrow \mathcal{I}^{\bullet}$$

is an injective resolution of $\mathcal{A}$ in the category of abelian sheaves over $\underline{\mathbf{X}}$ and Γ_{inv} denotes the invariant sections.

Example 3.11. Let $\underline{\mathbf{X}} = \mathbf{M} // \mathbf{G}$ be a global quotient by a finite group, let $\mathbf{V}$ be a $\mathbf{G}$ -equivariant complex vector bundle over $\mathbf{M}$, and denote its sheaf of sections by $\mathcal{O}(\mathbf{V})$. Since taking $\mathbf{G}$ -invariants is an exact functor on complex vector spaces (denoted $(-)^{\mathbf{G}}$), we have

$$\mathbf{H}^*(\underline{\mathbf{X}}, \mathcal{O}(\mathbf{V})) \cong \mathbf{H}^*(\mathbf{M}, \mathcal{O}(\mathbf{V}))^{\mathbf{G}}.$$

Definition 3.12. Let $\underline{X}$ be an orbifold, and let V be a complex vector bundle on $\underline{X}$. We define the topological Todd genus (also known as topological complex Euler characteristic) of $\underline{X}$ with coefficients in V by

$$\mathrm{Td}_{\mathrm{top}}(\underline{X}, V) := \mathrm{sdim} H^*(\underline{X}, \mathcal{O}(V)) = \sum_{\mathfrak{p}} (-1)^{\mathfrak{p}} H^{\mathfrak{p}}(\underline{X}, \mathcal{O}(V)).$$

If W is a complex vector bundle over $\Lambda \underline{X}$, then the orbifold Todd genus of $\underline{X}$ with coefficients in W is defined by

$$\mathrm{Td}_{\mathrm{orb}}(\underline{X}, W) := \mathrm{sdim} H^*(\Lambda \underline{X}, \mathcal{O}(W)).$$

Example 3.13. In the situation of Example 3.11, one has [AS68, pp.543-545]

$$\begin{aligned} \mathrm{Td}_{\mathrm{top}}(M//G; V) &= \mathrm{sdim} H^*(M; \mathcal{O}(V))^G \\ &= \frac{1}{|G|} \sum_{g \in G} \sum_{\mathfrak{p}} (-1)^{\mathfrak{p}} \mathrm{Trace}(g | H^{\mathfrak{p}}(M, \mathcal{O}(V))) \\ &= \frac{1}{|G|} \sum_{g \in G} \mathrm{Td} \left(M^g, \frac{V|_{M^g}}{\lambda_{-1}((N^g)^*)}(g) \right). \end{aligned}$$

Definition 3.14. We set

$$\mathrm{Td}_{\mathrm{top}} \left(\underline{X}; \sum_j V_j q^{\frac{j}{l}} \right) := \sum_j \mathrm{Td}_{\mathrm{top}}(\underline{X}; V_j) q^{\frac{j}{l}}.$$

Let $\underline{X}$ be an U^2 -orbifold, such that $\Lambda \underline{X}$ is connected. We set

$$\hat{A}_{\mathrm{orb}}^{\mathrm{string}}(\underline{X}) := q^{-\frac{d}{12} + \frac{\mathrm{age}(A^* \mu)}{2}} \cdot \mathrm{Td}_{\mathrm{orb}} \left(\underline{X}; \sqrt{\det T \underline{X}} \cdot \prod_{j \geq 1} \mathrm{Sym}_{q^{\frac{j}{l}}} (T \underline{X}^{\mathbb{C}})_{\zeta_l^j} \right).$$

If $\Lambda \underline{X}$ is not connected, one sums over its connected components.

In the case of a global quotient orbifold by a finite group, this definition specialized to Definition 3.4.

The author recently learned of the work of Dong, Liu and Ma, who defined Morita-invariant versions of elliptic genera in [DLM02]. It seems likely that their definitions are closely related to ours. A comparison to their work would be very interesting, because they are working with the index theorem on orbifolds. The authors of [dFLNU06] also started from an index theoretic discussion but do not formulate their results in a way that makes the Morita invariance of their arguments obvious.

4. LOOP SPACES OF SYMMETRIC POWERS

4.1. Actions of wreath products. Let G be a finite group. Recall that the wreath product $G \wr \Sigma_n$ has elements

$$(\underline{g}, \sigma) \in G^n \times \Sigma_n$$

which compose as follows:

$$(g_1, \dots, g_n, \sigma) \cdot (h_1, \dots, h_n, \tau) = (g_1 h_{\sigma^{-1}(1)}, \dots, g_n h_{\sigma^{-1}(n)}, \sigma \tau).$$

Let l be the order of $g_n \cdots g_1$ in G , and let σ be the n -cycle $(1 \dots n)$. Then the order of $(\underline{g}, \sigma)$ in $G \wr \Sigma_n$ equals ln .

An element $(\underline{h}, \tau)$ is in the centralizer of $(\underline{g}, \sigma)$ if and only if

$$\tau\sigma = \sigma\tau \quad \text{and} \quad \forall i : g_{\sigma(\tau(i))} h_{\tau(i)} = h_{\tau(\sigma(i))} g_{\sigma(i)}.$$

Let $(\underline{h}, \tau)$ be in the centralizer of $(\underline{g}, \sigma)$, let $(i_1, \dots, i_k)$ and $(j_1, \dots, j_k)$ be k -cycles in σ , and assume that $\tau(i_r) = j_{r+m}$. Then

$$\forall r \in \mathbb{Z}/k\mathbb{Z} : g_{j_r} h_{j_{r-1}} = h_{j_r} g_{i_{r-m}}.$$

In particular,

$$h_{j_k} g_{i_{k-m}} \cdots g_{i_{1-m}} = g_{j_k} \cdots g_{j_1} h_{j_k},$$

and the other h_{j_r} are determined by h_{j_k} and $\underline{g}$. Note that the element

$$h_{j_k} g_{i_{1-m}}^{-1} \cdots g_{i_k}^{-1} = g_{j_1}^{-1} \cdots g_{j_m}^{-1} h_{j_m}$$

conjugates $g_{j_k} \cdots g_{j_1}$ into $g_{i_k} \cdots g_{i_1}$.

Let M be a right G -manifold. Then $G \wr \Sigma_n$ acts on $\underline{x} \in M^n$ via

$$\underline{x} \cdot (\underline{g}, \sigma) = (x_{\sigma(1)} g_{\sigma(1)}, \dots, x_{\sigma(n)} g_{\sigma(n)}).$$

More generally, let $\mathbf{G} := (G_0, G_1, s, t, u, i)$ be an orbifold groupoid.

Definition 4.1. We define the n^{th} symmetric power of $\mathbf{G}$, denoted $\mathbf{G} \wr \Sigma_n$, to be the groupoid $\mathbf{G}^n \rtimes \Sigma_n$. Explicitly, $\mathbf{G} \wr \Sigma_n$ has objects G_0^n , morphisms $G_1^n \times \Sigma_n$. Its source sends the morphism $(\underline{g}, \sigma)$ to $(s(g_{\sigma(1)}), \dots, s(g_{\sigma(n)}))$, while its target sends it to $(t(g_1), \dots, t(g_n))$. The unit map sends the object $\underline{x}$ to $(u(x_1), \dots, u(x_n), 1)$. The multiplication of $(\underline{g}, \sigma)$ and $(\underline{h}, \tau)$, where $g_{\sigma(i)} = h_i$, is given by $(g_1 h_{\sigma^{-1}(1)}, \dots, g_n h_{\sigma^{-1}(n)})$; and the inverse of $(\underline{g}, \sigma)$ is given by $(i(g_{\sigma(1)}), \dots, i(g_{\sigma(n)}), \sigma^{-1})$.

One checks that this definition is Morita-invariant.

Example 4.2 (Symmetric powers of loop spaces). We have

$$(\mathcal{L}(M//G)) \wr \Sigma_n = \left(\coprod_{\underline{g} \in G^n} \mathcal{L}_{g_1} M \times \cdots \times \mathcal{L}_{g_n} M \right) // G \wr \Sigma_n,$$

where $(\underline{h}, \tau) \in G \wr \Sigma_n$ acts on the right-hand side by

$$(\gamma_1, \dots, \gamma_n) \mapsto (\gamma_{\sigma(1)} h_{\sigma(1)}, \dots, \gamma_{\sigma(n)} h_{\sigma(n)}).$$

Assume we have decompositions

$$\mathcal{V}|_X \cong \sum_{j \in \mathbb{Z}} (V_j \rtimes G) \mathbf{q}^{\frac{j}{N}} =: \mathbf{x} \quad \text{and} \quad \mathcal{W}|_Y \cong \sum_{l \in \mathbb{Z}} (W_l \rtimes H) \mathbf{q}^{\frac{l}{M}} =: \mathbf{y}.$$

Racall that the decomposition of the external product $(\mathcal{V} \times \mathcal{W})|_{X \times Y} = \mathcal{V}|_X \times \mathcal{W}|_Y$ is

$$\mathbf{x} \times \mathbf{y} := \text{pr}_X^*(\mathbf{x}) \oplus \text{pr}_Y^*(\mathbf{y}).$$

Now X^n is a $G \wr \Sigma_n$ -space, and $\mathcal{V}|_X \wr \Sigma_n$ becomes a $(\mathbb{S}^1 \times G) \wr \Sigma_n$ -bundle over it, which we may view as a $\mathbb{S}^1 \times (G \wr \Sigma_n)$ -bundle via the diagonal inclusion of $\mathbb{S}^1$ in $(\mathbb{S}^1)^n$. As a $\mathbb{S}^1 \times G^n$ -bundle,

$(\mathcal{V}|_X) \wr \Sigma_n$ is

$$(9) \quad d_n(x) := \sum_{j \in \mathbb{Z}^n} (V_{j_1} \times 0 \times \cdots \times 0) q^{\frac{j_1}{N}} \oplus \cdots \oplus (0 \times \cdots \times 0 \times V_{j_n}) q^{\frac{j_n}{N}} \\ \cong \sum_{j \in \mathbb{Z}^n} \sum_{k=0}^n W_{j,k} q^{\frac{j}{N}},$$

where $W_{j,k}$ consists of the $\binom{n}{k}$ summands in which V_j turns up exactly k times and the other summands are zero (in particular, $W_{j,0} = 0$). The symmetric group acts on $W_{j,k}$ by permutation of these summands.

4.2. Loop spaces of symmetric powers. We are now going to describe the loop space of $(M//G) \wr \Sigma_n$,

$$\mathcal{L}((M//G) \wr \Sigma_n) = \coprod_{[\underline{g}, \sigma]} \mathcal{L}_{(\underline{g}, \sigma)}(M^n) // C(\underline{g}, \sigma).$$

Let $[\sigma]$ correspond to the partition $n = \sum_k k N_k$, and assume that for each cycle of σ , we have fixed a “first” element i_1 , and thus a representation as

$$(i_1, \dots, i_k).$$

Theorem 4.3. *The component of $\mathcal{L}((M//G) \wr \Sigma_n)$ corresponding to $[\underline{g}, \sigma]$ is homeomorphic to*

$$(10) \quad \mathcal{L}_{(\underline{g}, \sigma)} M^n \cong \prod_k \prod_{(i_1, \dots, i_k)} {}_k \mathcal{L}_{g_{i_k} \cdots g_{i_1}} M,$$

where the second product runs over all k -cycles in σ . Let $(\underline{h}, \tau)$ be an element of $C(\underline{g}, \sigma)$ with $\tau(i_1) = j_{1+m}$ as in Section 4.1, and define its action on $\gamma \in {}_k \mathcal{L}_{g_{i_k} \cdots g_{i_1}} M$ by

$$(\gamma \cdot (\underline{h}, \tau))(t) := \gamma(m + t) h_{j_k} g_{i_1 - m}^{-1} \cdots g_{i_k}^{-1}.$$

Then $\gamma \cdot (\underline{h}, \tau)$ is a path in ${}_k \mathcal{L}_{g_{i_k} \cdots g_{i_1}} M$, and the homeomorphism is $C(\underline{g}, \sigma)$ -equivariant with respect to this action on the right-hand side. Moreover, it preserves the $\mathbb{S}^1$ -action by reparametrization of paths. There is a canonical map from the target of (10) with this action of $C(\underline{g}, \sigma)$ to

$$\prod_k ({}_k \mathcal{L}(M//G)) \wr \Sigma_{N_k}.$$

Denote its composite with (10) by $F_{(\underline{g}, \sigma)}$. As a map of orbifolds, $F_{(\underline{g}, \sigma)}$ is independent of the choices made.




FIGURE 3. An element of $\mathcal{L}_{((g_1, \dots, g_5); (135)(24))} M^5$ and its image in $\mathcal{L}_{g_5 g_3 g_1} M \times \mathcal{L}_{g_4 g_2} M$.

PROOF : The path

$$\gamma = (\gamma_1, \dots, \gamma_n)$$

in M^n is in $\mathcal{L}_{(\underline{g}, \sigma)} M^n$ if and only if

$$\forall i : \gamma_i(1) = \gamma_{\sigma(i)}(0) g_{\sigma(i)}.$$

I.e., if $\gamma(0) = (m_1, \dots, m_n)$, then for $1 \leq i \leq n$, γ_i is a path from m_i to $m_{\sigma(i)} g_{\sigma(i)}$. Let $(i_1, \dots, i_k)$ be a cycle in σ . The paths $\gamma_{i_k}, \gamma_{i_1} g_{i_1}$ to $\gamma_{i_{k-1}} g_{i_{k-1}} \cdots g_{i_1}$ compose to a path

$$\gamma := \gamma_{i_k} * \gamma_{i_1} g_{i_1} * \cdots * \gamma_{i_{k-1}} g_{i_{k-1}} \cdots g_{i_1}$$

in ${}_k \mathcal{L}_{g_{i_k} \cdots g_{i_1}} M$. This defines the homeomorphism of the proposition (see Figure 3). Its inverse sends a path $\gamma \in {}_k \mathcal{L}_{g_{i_k} \cdots g_{i_1}} M$ to $(\gamma_{i_r})_{r=1}^k$ with

$$\gamma_{i_k} = \gamma|_{[0,1]}$$

and

$$\gamma_{i_r}(t) := \gamma(r+t) g_{i_1}^{-1} \cdots g_{i_r}^{-1},$$

for $1 \leq r < k$. Let $(\underline{h}, \tau)$ be as in the proposition. Then the i_r^{th} path of $(\gamma_i)_{i=1}^n \cdot (\underline{h}, \tau)$ is $\gamma_{j_{r+m}} h_{j_{r+m}}$, and

$$\gamma_{j_{r+m}} h_{j_{r+m}} g_{i_r} \cdots g_{i_1} = \gamma_{j_{r+m}} g_{j_{r+m}} \cdots g_{j_1} (g_{j_1}^{-1} \cdots g_{j_m}^{-1} h_{j_m}).$$

It follows that the centralizer acts as claimed. Recall that

$${}_k \mathcal{L}(M//G) \cong \coprod_{[g]} {}_k \mathcal{L}_g M // C_g^k$$

and that if h is conjugate to g the groupoids ${}_k \mathcal{L}_g M // C_g^k$ and ${}_k \mathcal{L}_h M // C_h^k$ are related by an isomorphism which is canonical up to a natural transformation (i.e., it becomes canonical in the category of orbifolds). To be specific, fix representatives of the conjugacy classes of G , fix k , and fix an ordering $i_1, \dots, i_{N_k}$ of the k -cycles of σ . For $1 \leq \alpha \leq N_k$, set $g_\alpha := g_{i_{\alpha,k}} \cdots g_{i_{\alpha,1}}$, and let r_α be the representative of g_α . Then on the α^{th} factor, the canonical map of the proposition is represented by multiplication with a group element s_α which conjugates g_α into r_α (and hence maps ${}_k \mathcal{L}_{g_\alpha} M$ to ${}_k \mathcal{L}_{r_\alpha} M$). Let now $(\underline{h}, \tau) \in C_G(\underline{g}, \sigma)$ be as in the proposition. Then τ defines a permutation $\rho \in \Sigma_{N_k}$ of the set of k -cycles of σ . Let r be the representative of $[g_{j_k} \cdots g_{j_1}]$. Since $h_{j_k} \cdot g_{i_{1-m}}^{-1} \cdots g_{i_k}^{-1}$ conjugates $g_{j_k} \cdots g_{j_1}$ into $g_{i_k} \cdots g_{i_1}$, r is also the representative of the latter, and under our canonical map, right multiplication with $h_{j_k} \cdot g_{i_{1-m}}^{-1} \cdots g_{i_k}^{-1}$ translates into multiplication with an element of C_r combined with the permutation ρ of factors of the target

$$\prod_{\alpha} {}_k \mathcal{L}_{r_\alpha} M.$$

Rescaling a path by $t \mapsto t + m$ is the action of $m \in C_{r_\alpha}^k$.

Assume now that we have chosen a different first element, say i_r , of the cycle $(i_1, \dots, i_k)$. This leads to a different groupoid map $F'_{(\underline{g}, \sigma)}$. The factor of the target of $F_{(\underline{g}, \sigma)}$ corresponding to this cycle is $\mathcal{L}_{g_{i_k} \cdots g_{i_1}} M$, while that of $F'_{(\underline{g}, \sigma)}$ is $\mathcal{L}_{g_{i_{r-1}} \cdots g_{i_r}} M$. There is a natural isomorphism I between the two maps, sending $\gamma \in \mathcal{L}_{g_{i_k} \cdots g_{i_1}} M$ to

$$I(\gamma)(t) = \gamma(t+r) g_{i_1}^{-1} \cdots g_{i_{r-1}}^{-1}.$$

Hence F and F' define the same map of orbifolds. Similarly, changing the order of the k -cycles of σ translates into a permutation in Σ_{N_k} . $\square$

The orbifold loop space $\mathcal{L}((M//G) \wr \Sigma_n)$ can be viewed as the space of n strings moving in $M//G$, their order does not matter, they are either closed (in $M//G$) or joining together to form longer closed strings.

Example 4.4. Let $(\underline{h}, \tau)$ be equal to $(\underline{g}, \sigma) \in C_{(\underline{g}, \sigma)}$. Since $\sigma(i_1) = i_2$, and $g_{i_k} \cdot g_{i_k}^{-1} = 1$, its action on $\mathcal{L}_{g_{i_k} \dots g_{i_1}} M$ rotates the path $\gamma(t)$ to $\gamma(t+1)$. Hence $(\underline{g}, \sigma)$ maps to $1 \in C_r^k$ in the ${}_k\mathcal{L}_r M$ -factor of ${}_k\mathcal{L}(M//G)$ (see Figure 4).




FIGURE 4. The images in $\mathcal{L}_{g_3 g_2 g_1} M$ of an element of $\mathcal{L}_{(g_1, g_2, g_3; (123))} M^3$ and its counterpart under the action of $(g_1, g_2, g_3; (123))$ differ by rotation by 1.

In the case that G is the trivial group the discussion in [DMVV97] yields a formula summarizing the above description of $\mathcal{L}(M \wr \Sigma_n)$ for all n : write

$$\text{Sym}^\bullet(tM) := \coprod_{n \geq 0} M^n // \Sigma_n t^n$$

for the “total symmetric power” of M . Here t is a dummy variable.

Corollary 4.5. *We have*

$$\mathcal{L}(\text{Sym}^\bullet(tM)) \cong \prod_{k \geq 1} \text{Sym}^\bullet(({}_k\mathcal{L}(M) \rtimes (\mathbb{Z}/k)) t^k),$$

and the inertia orbifolds are

$$\Lambda(\text{Sym}^\bullet(tM)) := \prod_{n \geq 0} \Lambda(M \wr \Sigma_n) t^n \cong \prod_{k \geq 1} \text{Sym}^\bullet(t^k M \rtimes (\mathbb{Z}/k)),$$

where the action of the groups $\mathbb{Z}/k\mathbb{Z}$ on M is trivial.

For non-trivial G , these product formulas are proved in [Tam].

4.3. Loop bundles of symmetric products. In Section 5 we will define power operations in Tate K-theory. We want to define them in such a way that they are compatible with Thom classes in the sense of [tD68]. In order to understand the restrictions this condition places on our definitions, we will need to familiarize ourselves with the Fourier decomposition and the Thom-classes of loop bundles of symmetric powers of a vector bundle V .

Definition 4.6. Let $\mathcal{V}$ be an orbifold $\mathbb{S}^1$ -equivariant vector bundle over ${}_r\mathcal{LM} // G$. We write ${}_k\mathcal{V}$ for the orbifold $\mathbb{S}^1$ -equivariant vector bundle over $kr\mathcal{LM} // G$ obtained by rescaling the $\mathbb{R}/kl\mathbb{Z}$ action on $\mathcal{V}_g$ with $\frac{1}{k}$ to obtain an action of $\mathbb{R}/lkr\mathbb{Z}$.

Let $\mathcal{V}$ be as in the previous definition, and assume that its decomposition over the constant g -loops is

$$\mathcal{V}|_{M^g} \cong V_0 \bigoplus_{j \geq 1} V_j q^{\frac{j}{r+1}}.$$

Then the decomposition of ${}_k\mathcal{V}$ over M^g is

$${}_k\mathcal{V}|_{M^g} \cong V_0 \bigoplus_{j \geq 1} V_j q^{\frac{j}{kr+1}},$$

with the same V_j now viewed as $C_g^{\mathbb{R}/kr}$ representations, where C_g acts as before, and $\mathbb{R}/klr\mathbb{Z}$ acts with rotation number j . Hence the action of g on V_j equals that of $kr \in \mathbb{R}/klr\mathbb{Z}$, namely complex multiplication with ζ_l^j .

Definition 4.7. Let $\mathcal{V}$ be an orbifold $\mathbb{S}^1$ -equivariant vector bundle over $\mathcal{L}(M // G)$. We define the $\mathbb{S}^1$ -equivariant vector bundle $D_n(\mathcal{V})$ over $\mathcal{L}((M // G) \wr \Sigma_n)$ by

$$(D_n(\mathcal{V}))_{(\underline{g}, \sigma)} := F_{(\underline{g}, \sigma)}^* \left(\prod_k ({}_k\mathcal{V}) \wr \Sigma_{N_k} \right).$$

Example 4.8. In the case that $\mathcal{V} = \mathcal{L}(V)$ is the loop space of a bundle V over $M // G$, we have

$$D_n(\mathcal{V}) \cong \mathcal{L}((V // G) \wr \Sigma_n).$$

Let $(\underline{g}, \sigma)$ be as above, and let $f_{(\underline{g}, \sigma)} = F_{(\underline{g}, \sigma)}^{\mathbb{S}^1}$ be the restriction of $F_{(\underline{g}, \sigma)}$ to the constant loops.

Proposition 4.9. *Let the orbifold $\mathbb{S}^1$ -equivariant vector bundle $\mathcal{V}$ over $\mathcal{L}(M // G)$ have Fourier decomposition*

$$\mathfrak{x} := \mathcal{V}|_{\Lambda(M)} \cong \bigoplus_{j \in \mathbb{Z}} (V_j \wr G) q^{\frac{j}{|G|}}$$

over the G -space $\coprod_g M^g$. Then the restriction of $D_n(\mathcal{V})$ to $(M^n)^{(\underline{g}, \sigma)}$ has decomposition

$$f_{(\underline{g}, \sigma)}^* (d_{N_1}(1(\mathfrak{x})) \times d_{N_2}(2(\mathfrak{x})) \times \cdots \times d_{N_k}(k(\mathfrak{x})) \cdots)$$

over the $C_{G \wr \Sigma_n}(\underline{g}, \sigma)$ -space $(M^n)^{(\underline{g}, \sigma)}$. Here d_N is as in (9).

PROOF : By the definition of $f_{(\underline{g}, \sigma)}$, the diagram

$$\begin{array}{ccc} (M^n)^{(\underline{g}, \sigma)} & \xrightarrow{f} & \prod (\wedge(M//G)) \wr \Sigma_{N_k} \\ \downarrow i & & \downarrow \prod i_k \\ \mathcal{L}_{(\underline{g}, \sigma)} M^n & \xrightarrow{F} & \prod ({}_k \mathcal{L}(M//G)) \wr \Sigma_{N_k} \end{array}$$

commutes, where the products are over $k \in \mathbb{N}$, and i_k and i are the fixed point inclusions. Thus

$$\begin{aligned} (D_n \mathcal{V})|_{(M^n)^{(\underline{g}, \sigma)}} &= f_{(\underline{g}, \sigma)}^* \left(\prod_k i_k^* (({}_k \mathcal{V}) \wr \Sigma_{N_k}) \right) \\ &= f_{(\underline{g}, \sigma)}^* \left(\prod_k d_{N_k} (({}_k \mathcal{V})|_{\wedge(M//G)}) \right) \\ &= f_{(\underline{g}, \sigma)}^* \left(\prod_k d_{N_k} ({}_k \mathcal{X}) \right). \end{aligned}$$

Here $\prod$ stands for an external product. □

Definition 4.10. We define the map D_n in K_{Dev} by

$$\begin{aligned} D_n: K_{Dev, G}(M) &\rightarrow K_{Dev, G \wr \Sigma_n}(M^n) \\ x &\mapsto f_{(\underline{g}, \sigma)}^* (d_{N_1}(1(x)) \times d_{N_2}(2(x)) \times \cdots \times d_{N_k}(k(x)) \dots). \end{aligned}$$

4.4. Iterated symmetric powers. In order to prove that the definitions of section 5 satisfy the axioms for power operations, we will need to understand how our maps behave under iterated symmetric powers. Let

$$(\underline{\tau}, \sigma) = (\tau_1, \dots, \tau_n, \sigma) \in \Sigma_m \wr \Sigma_n,$$

and recall that for $1 \leq i \leq n$ and $1 \leq j \leq m$,

$$(\underline{\tau}, \sigma)(i, j) = (\sigma(i), \tau_{\sigma(i)}(j))$$

defines an action of $\Sigma_m \wr \Sigma_n$ on a set with nm elements, which is identified with $\{1, \dots, nm\}$ via $(i, j) \mapsto i + (j - 1)n$, inducing the inclusion

$$\iota: \Sigma_m \wr \Sigma_n \rightarrow \Sigma_{nm}.$$

If (i_r, j_r) is a cycle of $(\underline{\tau}, \sigma)$, then the i_r form a cycle of σ . Let σ be the cycle $(1 \dots n)$, and set

$$\tau = \tau_n \cdots \tau_1.$$

Then $(j_1 \dots j_k)$ is a k -cycle of τ , if and only if

$$((1, \tau_1(j_k))(2, \tau_2 \tau_1(j_k)) \dots (n, j_1)(1, \tau_1(j_1)) \dots (n, j_2)(1, \tau_1(j_2)) \dots (n, j_k))$$

is an nk -cycle of $(\underline{\tau}, \sigma)$.

Proposition 4.11. *Let*

$$(g_{i,j}, \tau_i, \sigma)_{i,j} \in (G \wr \Sigma_m) \wr \Sigma_n.$$

Then the following diagram commutes:

$$\begin{array}{ccc} \mathcal{L}_{(g_{i,j}, \tau_i, \sigma)_{i,j}}(M^m)^n & \longrightarrow & \prod_{\mathfrak{l}} \prod_{(i_1, \dots, i_{\mathfrak{l}})} \mathfrak{l} \mathcal{L}_{(g_j, \tau_j)} M^m \\ \parallel & & \downarrow \\ \mathcal{L}_{(g_{i,j}, \mathfrak{u}(\tau_i, \sigma))_{i,j}} M^{mn} & \longrightarrow & \prod_{\mathfrak{l}, (i_1, \dots, i_{\mathfrak{l}})} \prod_{\mathfrak{k}, (j_1, \dots, j_{\mathfrak{k}})} \mathfrak{k} \mathcal{L}_{g_{j_k} \dots g_{j_1}} M \end{array}$$

Here $(i_1, \dots, i_{\mathfrak{l}})$ runs over all $\mathfrak{l}$ -cycles of σ ,

$$(g_j, \tau)_j := (g_{i_{\mathfrak{l}}, j}, \tau_{i_{\mathfrak{l}}}) \cdots (g_{i_1, j}, \tau_{i_1}),$$

and the $(j_1, \dots, j_{\mathfrak{k}})$ run over all $\mathfrak{k}$ cycles of τ (hence the product in the lower right entry runs over all cycles of (τ, σ)). The upper map is $F_{((\underline{g}, \tau), \sigma)}$, the right vertical map is $\prod_{\mathfrak{l}, \mathfrak{i}} F_{(\underline{g}, \tau_{\mathfrak{i}})}$, the left is $\mathcal{L}\mathfrak{u}$ and the bottom is $F_{(\underline{g}, \mathfrak{u}(\tau, \sigma))}$.




FIGURE 5. An example with $\sigma = (12)$, $\tau_1 = (12)(3)$, and $\tau_2 = (123)$. Hence $\tau = (13)(2)$ and $g_1 = g_{21}g_{13}$ and $g_2 = g_{22}g_{11}$ and $g_3 = g_{23}g_{12}$.

PROOF : Without loss of generality, we may assume $\sigma = (1 \dots n)$. Then $F_{((\underline{g}, \tau), \sigma)}$ sends $(\gamma_{i,j})_{i,j}$ to $(\gamma_j)_{j=1}^m$ with

$$\gamma_j = \gamma_{n,j} * \gamma_{1, \tau_1(j)} g_{1, \tau_1(j)} * \cdots * \gamma_{n-1, \tau_{n-1} \cdots \tau_1(j)} g_{n-1, \tau_{n-1} \cdots \tau_1(j)} \cdots g_{1, \tau_1(j)}$$

(see Figure 5). Let

$$(g_1, \dots, g_m, \tau) := (g_{n,1}, \dots, g_{n,m}, \tau_n) \cdots (g_{1,1}, \dots, g_{1,m}, \tau_1).$$

Then $\tau = \tau_n \cdots \tau_1$, and

$$g_j = g_{n,j} g_{n-1, \tau_n^{-1}(j)} \cdots g_{1, \tau_2^{-1} \cdots \tau_n^{-1}(j)},$$

and γ_j is a path from $x_{n,j}$ to $x_{n, \tau(j)} g_{\tau(j)}$. Hence $(\gamma_j)_{j=1}^m$ is an element of

$$n \mathcal{L}_{(g_1, \dots, g_m, \tau)} M^m.$$

Let now $(j_1, \dots, j_{\mathfrak{k}})$ be a cycle of τ . Then the component of $f_{m,G}$ corresponding to this cycle sends (γ_j) to

$$\gamma = \gamma_{j_{\mathfrak{k}}} * \gamma_{j_1} g_{j_1} * \cdots * \gamma_{j_{\mathfrak{k}-1}} g_{j_{\mathfrak{k}-1}} \cdots g_{j_1}.$$

For $r \in \mathbb{Z}/k\mathbb{Z}$, we have

$$\gamma|_{[rn+i, rn+i+1]} = \gamma_{j_r} g_{j_r} \cdots g_{j_1}|_{[i, i+1]}.$$

Further,

$$\begin{aligned} \gamma_{j_r}|_{[i, i+1]} &= \gamma_{i, \tau_i \cdots \tau_1(j_r)} g_{i, \tau_i \cdots \tau_1(j_r)} g_{i-1, \tau_{i-1} \cdots \tau_1(j_r)} \cdots g_{\tau_1(j_r)} \\ &= \gamma_{(\underline{\tau}, \sigma)^{rn+i}(n, j_k)} g_{(\underline{\tau}, \sigma)^{rn+i-1}(n, j_k)} \cdots g_{(\underline{\tau}, \sigma)^{rn+1}(n, j_k)}, \end{aligned}$$

and

$$g_{j_s} = g_{(\underline{\tau}, \sigma)^{sn}(n, j_k)} \cdots g_{(\underline{\tau}, \sigma)^{(s-1)n+1}(n, j_k)}.$$

□

Corollary 4.12. *We have*

$$\text{res}|_t \circ D_{nm} \cong D_n \circ D_m.$$

5. POWER OPERATIONS

We will work with the equivariant notion of power operations spelled out in [Gan06]. Let E_G be a compatible family of equivariant cohomology theories with products. Let X be a G -space. Let $R_E(X)$ be the ring

$$R_E(X) = \bigoplus_{n \geq 0} E_{G \wr \Sigma_n}(X^n) t^n.$$

Here t is a dummy variable keeping track of the grading, and the multiplication of two elements of degree m and n respectively, is given by the external product, followed by the transfer

$$\text{ind}|_{\Sigma_m \times \Sigma_n}^{\Sigma_{m+n}}.$$

The ring $R_E(X)$ is the target of the total power operation

$$P = \sum_{n \geq 0} P_n t^n.$$

5.1. Atiyah power operations. The notion of Atiyah power operation in Tate K -theory is the continuation of Atiyah's definition of power operations in K -theory to that equivariant version of Tate K -theory, which is the straight-forward continuation of equivariant K -theory:

$$E_G(X) = K_G(X)[[q]].$$

Atiyah power operations provide the framework for studying the product formula for the topological Witten genus of an orbifold. Recall that Atiyah's power operations in real or complex K -theory send a class $x = [V]$ which is represented by a G -equivariant vector bundle over X , to the $G \wr \Sigma_n$ -equivariant bundle

$$P_n(V) := V^{\boxtimes n},$$

over X^n (here Σ_n acts with a sign on $V^{\boxtimes n}$).

Definition 5.1. We define the total Atiyah power operation on K_{Tate} by

$$\begin{aligned} P^{\text{top}}: K_G(X)[[q]] &\rightarrow \bigoplus_{n \geq 0} K_{G \wr \Sigma_n}(X^n)[[q]] t^n \\ x_j q^j &\mapsto \sum_{n \geq 0} x_j^{\otimes n} q^{jn} t^n, \text{ and} \\ \sum_{j \geq 0} x_j q^j &\mapsto \prod_{j \geq 0} P^{\text{top}}(x_j q^j). \end{aligned}$$

The coefficient of t^n is called the n^{th} Atiyah power operation, it is given by

$$P_n^{\text{top}}(x) = \sum_{\underline{i}} P_{\underline{i}}(x),$$

where $\underline{i} = (i_j)_{j \geq 0}$ runs over all sequences of natural numbers whose elements add up to n , and

$$P_{\underline{i}}(x) = \prod_j P_{i_j}(x_j) q^{i_j j}.$$

Note that for each $\underline{i}$, this product has only finitely many non-trivial factors. It is taken in $K_G(X)$ and hence involves the induced representations

$$\text{ind } |\underline{i}|_i^n := \text{ind } |\Sigma_{i_1} \times \dots \times \Sigma_{i_j} \times \dots$$

The transfer $\text{ind } |\underline{i}|_i^n$ adds all possible shuffles of the partition $\underline{i}$. More precisely, we have

$$(11) \quad P_n^{\text{top}}(x) = \sum_{j \in \mathbb{N}^n} x_{j_1} q^{j_1} \boxtimes \dots \boxtimes x_{j_n} q^{j_n},$$

and Σ_n acts (with a sign) by permutation of the summands and the factors therein. The infinite sum P_n^{top} is a well defined formal power series in q .

Proposition 5.2. *The P_n^{top} satisfy the axioms of [Gan06, Def.4.3].*

PROOF : Part (a) is clear. Part (b) and (c) follow from Equation (11). Let $x \in K_G(X)[[q]]$ and $y \in K_H(Y)[[q]]$. Then the external product of x and y is

$$x \boxtimes y = \left(\sum_i x_i q^i \right) \boxtimes \left(\sum_j y_j q^j \right) = \sum_{i,j} (x_i \boxtimes y_j) q^{i+j},$$

and

$$P_n(x \boxtimes y) = \sum_{i,j} (x_{i_1} \boxtimes y_{j_1}) q^{i_1+j_1} \boxtimes \dots \boxtimes (x_{i_n} \boxtimes y_{j_n}) q^{i_n+j_n},$$

Under the isomorphism

$$(X \wedge Y)^n \cong X^n \wedge Y^n,$$

this is identified with

$$\text{res } |_{(G \times H) \wr \Sigma_n}^{(G \wr \Sigma_n) \times (H \wr \Sigma_n)} P_n(x) \boxtimes P_n(y).$$

□

Let V be a G -equivariant $\text{Spin}(2d)$ vector bundle, and set

$$u_G^{\text{top}}(V) := \cdot u_G^{\hat{\Lambda}}(V) \cdot \prod_{j=1}^{\infty} \Lambda_{-q^j} V_{\mathbb{C}}.$$

This is a Thom class in $K_G(X^V)[[q]]$, and hence, by Proposition A.4, we have a map

$$\phi_G^{\text{top}}: M\text{Spin}P_G \rightarrow K_G[[q]].$$

Similarly, replacing $\hat{A}$ by Td , we obtain Thom classes for complex vector bundles, and a map

$$MUP_G \rightarrow K_G[[q]].$$

Proposition 5.3. *The map ϕ_G^{top} is H_∞^4 (H_∞^2 in the complex case).*

PROOF : By [Gan06, 4.6],[tD68, (A4)], it suffices to show that

$$u_{G|\Sigma_n}^{\text{top}}(V^n) = P_n(u_G^{\text{top}}(V))$$

for any $\text{Spin}(4k)$ -bundle V . This follows from Axiom (d), $\dim V^n = 4nk$, the known analogue for $u^{\hat{A}}$ and u^{Td} :

$$u_{G|\Sigma_n}^{\hat{A}}(V^n) = P_n(u_G^{\hat{A}}(V))$$

[BMMS86], and the fact that

$$\Lambda_t(V^n) \cong (\Lambda_t V)^{\boxtimes n},$$

where Σ_n acts by permuting the factors inside on the left-hand side, and by permuting the factors outside (with a sign) on the right-hand side. $\square$

Definition 5.4. We define the topological Witten genus of a G -manifold M by

$$\phi_{\text{top}}(M//G) := \frac{1}{|G|} \sum_{g \in G} \text{Trace}(g | \phi_G^{\text{top}}(M)),$$

where ϕ stands for $\hat{A}$ or Td .

Definition 5.5. We define the (topological) total symmetric power operation $\text{Sym}_t^{\text{top}}$ on K_{Tate} to be the composite

$$\text{Sym}_t^{\text{top}}: K_G(X)[[q]] \rightarrow \bigoplus_{n \geq 0} K_{G|\Sigma_n}(X^n)[[q]]t^n \rightarrow \bigoplus_{n \geq 0} K_{G \times \Sigma_n}(X)[[q]]t^n \rightarrow K_G(X)[[q]][[t]],$$

where the first map is P^{top} , and the second map is restriction along the diagonals of X^n and G^n . The last map ε is (in degree n) given by

$$\varepsilon_n = \text{id} \otimes \langle -, 1 \rangle_{\Sigma_n}: K_G(X) \otimes R(\Sigma_n) \rightarrow K_G(X).$$

Note that the total power operation takes sums to products, and that ε is a ring map. Hence S_t is exponential. In the special case that X is a point and G is the trivial group, we obtain the following corollary of Proposition 5.3:

Corollary 5.6 (product formula for the topological Witten genus). *If M is a $\text{Spin}(4k)$ or complex manifold with Witten genus $\phi(M) = \sum_j c(j)q^j$, then*

$$\sum_{n \geq 0} \phi_{\text{top}}(M^n//\Sigma_n)t^n = \prod_j \left(\frac{1}{1 - tq^j} \right)^{c(j)}.$$

PROOF : Since ϕ^{top} preserves power operations, we have

$$\sum_{n \geq 0} \phi_{\Sigma_n}^{\text{top}}(M^n) t^n = P^{\text{top}}(\phi(M)).$$

Applying ε to both sides, we get

$$\sum_{n \geq 0} \phi_{\text{top}}(M^n // \Sigma_n) t^n = \text{Sym}_t^{\text{top}}(\phi(M)).$$

Since $\text{Sym}_t^{\text{top}}$ is exponential, it suffices to observe that $\text{Sym}_t^{\text{top}}(q^j) = \frac{1}{1-tq^j}$ to complete the proof. $\square$

5.2. Stringy power operations.

Definition 5.7. Let $\kappa(-)$ be the ring map

$$\begin{aligned} \kappa(-): K_{\text{Dev}, G, r} &\rightarrow K_{\text{Dev}, G, rk} \\ \sum V_j q^{\frac{j}{r|g|}} &\mapsto \sum V_j q^{\frac{j}{rk|g|}}. \end{aligned}$$

Here the action of C_g on V_j remains unchanged, while $1 \in \mathbb{Z}/rk|g|\mathbb{Z}$ acts as $\zeta_{rk|g|}^j$ on the j^{th} coefficient of the $[g]^{\text{th}}$ summand on the right-hand side.

In the following, for $x \in K_{\text{Dev}, G, \kappa}(X)$, by $P_n^{\text{top}}(x)$ we will mean the image of x under the composite

$$K_{\text{Dev}, G, \kappa}(X) \rightarrow K_{G \times \mathbb{Z}/kL}(\coprod_g X^g) \llbracket q^{\frac{1}{kL}} \rrbracket \rightarrow K_{(G \times \mathbb{Z}/kL) \wr \Sigma_n}((\coprod_g X^g)^n) \llbracket q^{\frac{1}{kL}} \rrbracket.$$

Here L is the order of G .

Definition 5.8. Let $x \in K_{\text{Dev}, G}(X)$. We define

$$P_n^{\text{string}}(x) \in K_{\text{Dev}, G \wr \Sigma_n}(X^n)$$

by

$$(P_n^{\text{string}}(x))_{(\underline{g}, \sigma)} := f_{(\underline{g}, \sigma)}^* (P_{N_1}^{\text{top}}(x_1) \boxtimes \cdots \boxtimes P_{N_k}^{\text{top}}(x_k) \boxtimes \cdots).$$

We need to verify that P_n^{string} indeed takes values in $K_{\text{Dev}, G \wr \Sigma_n}(X^n)$. Firstly, we have

$$\begin{aligned} P_{N_k}^{\text{top}}(x_k) &\in K_{(G \times \mathbb{Z}/kL) \wr \Sigma_{N_k}}((\coprod_{h \in G} X^h)^{N_k}) \llbracket q^{\frac{1}{kL}} \rrbracket \\ &\cong \bigoplus_{[\underline{h}]} K_{\text{Stab}(\underline{h})}(\prod_{i=1}^{N_k} X^{h_i}) \llbracket q^{\frac{1}{kL}} \rrbracket, \end{aligned}$$

where the direct sum is over the orbits of the action of $(G \times \mathbb{Z}/kL) \wr \Sigma_{N_k}$ on G^{N_k} . In fact, the $\underline{h}^{\text{th}}$ summand of $P_{N_k}^{\text{top}}(x_k)$ is a power series in $q^{\frac{1}{k|\underline{h}|}}$, since it is obtained by multiplying power series in $q^{\frac{1}{k|\underline{h}_i|}}$ for $1 \leq i \leq N_k$. Further, recall that

$$(X^n)^{(\underline{g}, \sigma)} \cong \prod_{k \geq 1} \prod_{(i_1, \dots, i_k)} X^{g_{i_k} \cdots g_{i_1}},$$

27

where the second product runs over the k -cycles of σ . Hence, for fixed k , the target of the k^{th} factor of $f_{(\underline{g}, \sigma)}$ is $\prod_{i=1}^{N_k} X^{h_i}$, where i runs over the k -cycles of σ , and $h_i = g_{i_k} \cdots g_{i_1}$. By Example 4.4, $(\underline{g}, \sigma)$ acts by multiplication with $\zeta_{k|h_i|}^j$ on the coefficient of $q^{\frac{j}{k|h_i|}}$ of $kx|_{X^{h_i}}$. Since $|(\underline{g}, \sigma)|$ is the smallest common multiple of all $k \cdot |g_{i_k} \cdots g_{i_1}|$ such that $k \in \mathbb{N}$ and $(i_1 \dots i_k)$ is a k -cycle of σ , it follows that $(p_n^{\text{string}}(x))_{(\underline{g}, \sigma)}$ is a power series in $q^{\frac{1}{|(\underline{g}, \sigma)|}}$, where $(\underline{g}, \sigma)$ acts by multiplication with $\zeta_{|(\underline{g}, \sigma)|}^j$ on the coefficient of $q^{\frac{j}{|(\underline{g}, \sigma)|}}$.

Theorem 5.9. *The p_n^{string} satisfy the axioms of [Gan06, Def.4.3].*

PROOF : Axiom (a) is clear. For Axiom (b), let $(\sigma, \tau) \in \Sigma_m \times \Sigma_n$. Then the set of k -cycles of the element $(\sigma, \tau) \in \Sigma_{m+n}$ is identified with the disjoint union of the sets of k -cycles of σ and τ , and thus

$$\begin{aligned} (p_{n+m}^{\text{string}}(x))_{(\underline{g}, (\sigma, \tau))} &= (p_m^{\text{string}})_{(g_1, \dots, g_m, \sigma)} \boxtimes (p_n^{\text{string}})_{(g_{m+1}, \dots, g_{m+n}, \tau)} \\ &= (p_m^{\text{string}} \boxtimes p_n^{\text{string}})_{(\underline{g}, (\sigma, \tau))}. \end{aligned}$$

Axiom (c) follows from Proposition 4.11: With the notation of that Proposition, and omitting the equivariant information, we have

$$\begin{aligned} (p_n^{\text{string}}(p_m^{\text{string}}(x)))_{(g_{i,j}, \tau_i, \sigma)} &= \prod_{l, (i_1, \dots, i_l)} l \left(\prod_{k, (j_1, \dots, j_k)} kx_{g_{j_k} \cdots g_{j_1}} \right) \\ &= p_n^{\text{string}}(x)_{(g_{i,j}, \tau_i, \sigma)}, \end{aligned}$$

Here all the products denote the external tensor product $\boxtimes$. For the non-equivariant information it makes no difference whether $l(-)$ is inside or outside the product, and when restricting along the inclusion of centralizers

$$C_{G(\Sigma_m \wr \Sigma_n)}(g_{i,j}, \tau_i, \sigma) \subseteq C_{G(\Sigma_{mn})}(g_{i,j}, \tau_i, \sigma)$$

we restrict that part along the inclusion $\mathbb{Z}/l\mathbb{Z} \subseteq (\mathbb{Z}/lk\mathbb{Z})^{N_k}$. To prove Axiom (d), let $x \in K_{\text{Dev}, G}(X)$ and $y \in K_{\text{Dev}, H}(Y)$. Then $x \boxtimes y$ is the element of $K_{\text{Dev}, G \times H}$ with $(x \boxtimes y)_{[g, h]} = x_{[g]} \boxtimes y_{[h]}$, and the statement follows from the analogous statement for the $P_{N_k}^{\text{top}}$. $\square$

In this section, we will work with the following variant of the stringy equivariant Witten genus:

Definition 5.10. We define

$$\text{Td}_G^{\text{string}}: \text{MU}_G \rightarrow K_{\text{Dev}}$$

to be the map of spectra corresponding to the Thom class

$$u_{\text{Td}, G}^{\text{string}}(V)_{[g]} := u_{C_g}^{\text{Td}}(V^g) \cdot \prod_{j \geq 1} \wedge_{-1} \left((V_C)_{\zeta_l^j} q^{\frac{j}{l}} \right).$$

The computation of $\text{Td}_{[g]}^{\text{string}}(M)(h)$ is the same as that for $\hat{A}^{\text{string}}$ with the terms

$$q^{-\frac{d}{12} + \frac{\text{age}(g)}{2}} e^{i\pi \text{age}(h)} \prod e^{\tilde{x}_i}$$

missing. For SU-bundles, we have $\sum \tilde{x}_i = 0$, so that the missing factors are just a renormalization constant.

Theorem 5.11. *The map $\mathrm{Td}_G^{\mathrm{string}}$ is H_∞^2 .*

PROOF : We write $\mathbf{u}_G^{\mathrm{string}}$ for $\mathbf{u}_{\mathrm{Td},G}^{\mathrm{string}}$. We need to prove

$$\mathbf{u}_{G|\Sigma_n}^{\mathrm{string}}(V^n) = \mathbf{p}_n^{\mathrm{string}}(\mathbf{u}_G^{\mathrm{string}}(V)).$$

Let $(\underline{g}, \sigma) \in G \wr \Sigma_n$. Then

$$V^n|_{(X^n)^{(\underline{g}, \sigma)}} \cong \prod_k \prod_{\underline{i}} V^{\oplus k}|_{X^{g_{i_k} \cdots g_{i_1}}},$$

where the product signs denote external direct sums and the second one runs over the k -cycles of σ . On the $\underline{i}^{\mathrm{th}}$ summand of this, $(\underline{g}, \sigma)$ acts by

$$(v_1, \dots, v_k) \mapsto (v_2 g_{i_2}, \dots, v_k g_{i_k}, v_1 g_{i_1}).$$

Hence ζ is an eigenvalue of this action with eigenvector $(v_1, \dots, v_k)$ if and only if ζ^k is an eigenvalue of $g_{i_k} \cdots g_{i_1}$ with eigenvector v_k , and $v_j = v_{j+1} g_{i_{j+1}}$ for $1 \leq j \leq k$. Thus

$$(V^{\oplus k}|_{X^{g_{i_k} \cdots g_{i_1}}})_{\zeta} \cong (V|_{X^{g_{i_k} \cdots g_{i_1}}})_{\zeta^k},$$

and

$$\left(V^n|_{(X^n)^{(\underline{g}, \sigma)}} \right)_{\zeta} \cong \prod_k \prod_{\underline{i}} (V|_{X^{g_{i_k} \cdots g_{i_1}}})_{\zeta^k}.$$

Now Definition 5.10 gives

$$\begin{aligned} \mathbf{u}_{(\underline{g}, \sigma), \alpha}^{\mathrm{string}}(V^n) &= \mathbf{u}_{C_{(\underline{g}, \sigma)}}(V_1) \cdot \prod_{j=1}^{\infty} \wedge_{-q^{\frac{j}{|\underline{g}, \sigma|}}} \left((V_{\mathbb{C}})_{\zeta_{|\underline{g}, \sigma|}^j} \right) \\ &= \prod_{k, \underline{i}} \left(\mathbf{u}_{C_{g_{\underline{i}}}}(V_1) \cdot \prod_{j=1}^{\infty} \wedge_{-q^{\frac{j}{k|\underline{g}_{\underline{i}}|}}} \left((V_{\mathbb{C}})_{\zeta_{|\underline{g}_{\underline{i}}|}^j} \right) \right), \end{aligned}$$

where $g_{\underline{i}} = g_{i_k} \cdots g_{i_1}$, in the first line, V and $V_{\mathbb{C}}$ get restricted to $(X^n)^{(\underline{g}, \sigma)}$ and in the second line, they get restricted to $X^{g_{i_k} \cdots g_{i_1}}$. $\square$

5.3. Hecke operators. Recall from [Gan06] that in a cohomology theory with power operations and a level 2 Hopkins-Kuhn-Ravenel theory, one defines Hecke operators by

$$(12) \quad T_n(x) := \frac{1}{n} \sum_{\alpha \in A} \psi_{\alpha}(x).$$

Here A is a system of representatives of those conjugacy classes of pairs of commuting elements¹ (σ, τ) of Σ_n , with the property that the subgroup of Σ_n generated by σ and τ acts transitively on $\{1, \dots, n\}$. For such a representative $\alpha \in A$, the operation ψ_{α} is defined by

$$\psi_{\alpha}(x) := \mathrm{eval}_{\alpha}(\mathrm{res}|_{\delta} P_n(x)),$$

¹In this context, “conjugacy class” refers to simultaneous conjugation.

where δ denotes the inclusion of the groupoid $X \rtimes (G \times \Sigma_n)$ into $(X \rtimes G) \wr \Sigma_n$, which is induced by the diagonal inclusion of $X \rtimes G$ into $(X \rtimes G)^n$. (On the source of δ , Σ_n acts trivially on the G -space X .)

In this section, we are going to compute T_n . We start by choosing a system of representatives A as in (12). In order for (σ, τ) to act transitively, any two cycles of σ must have equal length. Let c_1 be a cycle of σ , w.l.o.g. say $c_1 = (1, \dots, k)$. Let c_2 be the k -cycle of σ starting with $\tau(1)$. Noting that c_2 cannot equal c_1 , because of the transitivity of the action, we may assume $c_2 = (k+1, \dots, 2k)$. Repeating this argument, we arrive at N k -cycles $c_1, \dots, c_N$ of σ , where $n = Nk$, and for $1 \leq j < N$, τ maps the first element of c_j to the first element of c_{j+1} . For definiteness, we let $c_j = ((j-1)k+1, \dots, jk)$. Now τ has to map the first element of c_N to an element of c_1 , and each of the elements of c_1 is possible. The conjugacy class of (σ, τ) is uniquely determined by k and $1 + m := \tau((N-1)k+1) \in \{1, \dots, k\}$.

Fix N and k such that $Nk = n$, and let σ be the element of Σ_n with N k -cycles as above. Let $g \in G$ and $\underline{g} := \delta(g) = (g, \dots, g) \in G^n$. Then

$$P_{[\underline{g}, \sigma]}^{\text{string}}(x) = f_{(\underline{g}, \sigma)}^* P_N^{\text{top}}(kx|_{X^{g^k}}).$$

Consider $(h, \tau) \in C_G(g) \times C_{\Sigma_n}(\sigma)$, with τ as above. In order to determine its action on $(X^{g^k})^N$, we note that we can simplify the discussion in the proof of Theorem 4.3 by choosing all the representatives r_α to equal g^k . Then (h, τ) acts as

$$(h, \dots, h, (hg^{-m}, m); C) \in C_g^k \wr \Sigma_N$$

on $(X^{g^k})^N$, where $C = (1, \dots, N)$.

Assume first that $m = 0$. Then P_N^{top} composed with $\text{Trace}(-, \tau)$ is the N^{th} Adams operator ψ_N . The set of τ -fixed points of $(X^{g^k})^N$ is the image of the diagonal map

$$\delta': X^{g^k} \rightarrow (X^{g^k})^N.$$

Hence the $[g]$ -component of $\psi_{\sigma, \tau}(x)$ is

$$(13) \quad (\psi_{\sigma, \tau}(x)) [g] = i^*(\delta')^*(\psi_N(kx|_{X^{g^k}})),$$

where

$$i: X^g \supset C_g \rightarrow X^{g^k} \supset C_{g^k}^k$$

is the inclusion. On coefficients, we get the following formula for characters:

$$(P_n^{\text{string}}(x))(\underline{g}, \sigma; \underline{h}, \tau) = \left(\sum_{j \geq 0} V_j q^{\frac{Nj}{kl}}(g^k, h^N) \right),$$

where l is the order of g , and $x = \sum_{j \geq 0} V_j q^{\frac{j}{l}}$. Let now m be arbitrary. Then $(h, m) \in C_{g^k}^k$ acts as $h \cdot \zeta_{lk}^{mj}$ on V_j . Non-equivariantly, (13) is still true, however, now the action of h on $\psi_N(kx|_{X^{g^k}})$ is twisted with $g^{-m} \zeta_{lk}^{mj}$. On coefficients, we get

$$(P_n^{\text{string}}(x))(\underline{g}, \sigma; \underline{h}, \tau) = \left(\sum_{j \geq 0} V_j q^{\frac{Nj}{kl}} \zeta_{lk}^{jm}(g^k, g^{-m} h^N) \right),$$

where x and l are as above.

Remark 5.12. Viewing $\mathfrak{x}(\mathfrak{g}, \mathfrak{h})$ as the $\mathfrak{q}$ expansion of a function $\mathfrak{x}(\mathfrak{g}, \mathfrak{h}; z)$, where z is in the upper half-plane, and $\mathfrak{q} = e^{2\pi iz}$, we get (on coefficients)

$$(T_n(\mathfrak{x}))(\mathfrak{g}, \mathfrak{h}; z) = \frac{1}{n} \sum_{\substack{Nk=n \\ 0 \leq m < k}} \mathfrak{x}\left(\mathfrak{g}^k, \mathfrak{g}^{-m}\mathfrak{h}^N, \frac{Nz + m}{k}\right).$$

This is exactly the formula for the twisted Hecke operators of generalized Moonshine in [Gan]. It follows from the way the twisted Hecke operators are defined in [ibid.] (namely using isogenies of elliptic curves) that our $\mathbf{P}_n^{\text{string}}$ are elliptic. Let G be the trivial group. Using the theory of isogenies on the Tate-curve, Matthew Ando has defined power operations on K_{Tate} in [And00]. In this case, our definition specializes to his. For $G = 1$ and X a point, the Hecke operators are the usual ones acting on the $\mathfrak{q}$ -expansions of modular forms.

As we noted above, our geometric picture only sees one of the two circles of the elliptic curve. The case where $k = n$ and $N = 1$ corresponds to replacing this circle by one of length n and pulling back along an n -fold covering map from this long circle to the short one. The analogous power operation for the circle we cannot see ($k = 1$ and $N = n$), is the n^{th} Adams operation.

5.4. The DMVV-formula, Borchers products and replicability. We recall from [Gan06] that in a cohomology theory with power operations and a level 2 Hopkins-Kuhn-Ravenel theory, the n^{th} (stringy) symmetric power is defined by

$$\text{sym}_{\text{string}}^n(\mathfrak{x}) = \frac{1}{n!} \sum_{\alpha} \psi_{\alpha}(\mathfrak{x}),$$

A similar argument to that of Section 3.4 shows that the symmetric powers take values in $K_{\text{Dev}, G}(X)$. The total (stringy) symmetric power is defined as

$$\text{Sym}_{\mathfrak{t}}^{\text{string}}(\mathfrak{x}) := \sum \text{sym}_{\text{string}}^n(\mathfrak{x}) \mathfrak{t}^n.$$

With these definitions, the generating function argument in [Gan06, 9.2] has the following corollary:

Corollary 5.13. *On the level of cohomology operations, we have*

$$\text{Sym}_{\mathfrak{t}}^{\text{string}}(\mathfrak{x}) = \exp \left(\sum_{m \geq 1} T_m(\mathfrak{x}) \mathfrak{t}^m \right).$$

In terms of elliptic curves, this generating function can be interpreted as follows: the Hecke operators average over the pullback along a system of representatives for isogenies onto the elliptic curve. The n^{th} symmetric power averages over all possible n -fold coverings of the elliptic curve. Each covering is made up from its connected components, and those are isogenies. For a detailed discussion of this picture, we refer the reader to [Gan].

Together with Theorem 5.11, and applied to the case where X is a point, Corollary 5.13 becomes the Dijkgraaf-Moore-Verlinde-Verlinde formula:

Corollary 5.14. *Let M be a compact closed complex manifold. Then*

$$\sum_{n \geq 0} \text{Td}_{\text{orb}}^{\text{string}}(M^n) \mathfrak{t}^n = \exp \left(\sum_{m \geq 1} T_m(\text{Td}^{\text{string}}(M)) \mathfrak{t}^m \right).$$

The right-hand side of the DMVV-formula can be rewritten as a Borcherds product:

$$\prod_{i,j} \left(\frac{1}{1 - q^i t^j} \right)^{c(ij)},$$

where $\phi(M) = \sum c(j)q^j$. The best known example of a product formula is the case where the $c(i)$ are the coefficients of the q -expansion of the modular function $j - 744$. In this case, the product is equal to the inverse of

$$t(j(t) - j(q)).$$

In other words, all the mixed terms in t and q in the product $t^{-1} \prod (1 - q^i t^j)^{c(ij)}$ are zero. This property encodes the replicability of the function $j - 744$. Similar identities hold for the other Moonshine functions.

Note that $j(q) - 744$ equals the Witten genus of the Witten manifold M constructed in [MH02].

One rather peculiar property of these kind of product formulas is the change of the role of t from a dummy variable to a variable which plays the same role as q . In our context, both the variables t and q are counting winding numbers, which makes it less surprising that they should play a similar role. Indeed, the symmetry of their roles is clarified by the following proposition:

Proposition 5.15. *Let G be the trivial group, and view $x \in K(X)$ as an element of $K_{\text{Tate}}(X)$. Then the total symmetric power $\text{Sym}_q^{\text{string}}(x)$ is Witten's exponential characteristic class*

$$\text{Sym}_q^{\text{string}}(x) = \bigotimes_{k \geq 1} \text{Sym}_{q^k}(x),$$

where Sym_q stands for the total symmetric power in K -theory.

PROOF : Note that ${}_k x = \text{res}_{\mathbb{Z}/k}^1 x$. Hence, for every natural number N , we have

$$\langle P_N^{\text{top}}({}_k x), 1 \rangle_{\mathbb{Z}/k \wr \Sigma_N} = \langle P_N^{\text{top}}(x), 1 \rangle_{\Sigma_N} = \text{Sym}^N(x).$$

The coefficient of q^n in $\text{Sym}_q^{\text{string}}(x)$ is

$$\begin{aligned} \text{sym}_{\text{string}}^n(x) &= \frac{1}{|\Sigma_n|} \sum_{\sigma \tau = \tau \sigma} P_n^{\text{string}}(x)|_{\delta_n}(\sigma, \tau) \\ &= \sum_{[\sigma]} \frac{1}{|C_\sigma|} \sum_{\tau \in C_\sigma} \left(\bigotimes P_{N_k}^{\text{top}}({}_k x)|_{\delta_{N_k}} \right)(\tau), \end{aligned}$$

where $[\sigma]$ corresponds to the partition $n = \sum k N_k$, so that $C_\sigma \cong \prod_{k \geq 1} (\mathbb{Z}/k) \wr \Sigma_{N_k}$. (Since we are pulling back along the diagonal, the tensor product is an internal tensor product.) The expression becomes

$$\sum_{n = \sum N_k} \bigotimes_{k \geq 1} \left(\frac{1}{|k^{N_k} N_k!|} \sum_{\tau_k} \left(P_{N_k}({}_k x)|_{\delta_{N_k}} \right)(\tau_k) \right) = \sum_{n = \sum k N_k} \bigotimes_{k \geq 1} \text{Sym}^{N_k}(x).$$

(Here $\tau_k \in \mathbb{Z}/k \wr \Sigma_{N_k}$.) This is the coefficient of q^n in $\bigotimes_{k \geq 1} \text{Sym}_{q^k}(x)$. □

5.4.1. *Replicability.* Let $F(q)$ be a power series with coefficients in $\mathbf{R}(G)$ which is of the form

$$F(q) = q^{-1} + a_1 q + a_2 q^2 + \dots$$

From the Moonshine literature, such F are known as McKay-Thompson series. We recall² that the Faber polynomials $\Phi_{n,F}$ of F are defined by

$$-\sum_{n=1}^{\infty} \Phi_{n,F}(w) t^n = \log(t(F(t) - w)).$$

Hence $\Phi_{n,F}$ is a polynomial in w of degree n , which depends on the first n coefficients of F and is uniquely characterized by the fact that it is of the form

$$\Phi_{n,F}(F(q)) = q^{-n} + b_1 q + b_2 q^2 + \dots$$

Viewing $F(q)$ as an element in (the $[1]$ -component of) $q^{-1}K_{\text{Dev},G}(\text{pt})$, we arrive at the following definition:

Definition 5.16. Let F be a McKay-Thompson series. We write $F^{(a)}$ for the a^{th} Adams operator applied to F . We call F *replicable*, if for every natural number n , we have

$$\Phi_{n,F}(F(q)) = \sum_{\substack{a+d=n \\ 0 \leq b < d}} F^{(a)}\left(\frac{a\tau + b}{d}\right).$$

Here $q = e^{2\pi i \tau}$.

This appears to be the right notion of replicability of McKay-Thompson series, it is the one that turns up in [Bor92]. Note that the right-hand side of the equation in the definition equals $n \cdot T_n(F(q))$, where T_n is the Hecke operator computed in Section 5.3. It follows immediately from the definitions and from Corollary 5.13 that a McKay-Thompson series F is replicable if and only if it satisfies the following identity:

$$\begin{aligned} F(t) - F(q) &= t^{-1} \cdot \text{Sym}_t^{\text{string}}(-F(q)) \\ &= t^{-1} \cdot \Lambda_{-t}^{\text{string}}(F(q)), \end{aligned}$$

where $\Lambda_{-t}^{\text{string}}(x)$ is defined as the multiplicative inverse of $\text{Sym}_t^{\text{string}}(x)$. For $F(q)$ as above, $\Lambda_{-t}^{\text{string}}(F(q))$ can be written as

$$\Lambda_{-t}^{\text{string}}(F(q)) = \Lambda_{-1} \left(\sum a_{m,n} q^m t^n \right).$$

This is the form in which it appears in [Bor92, p.410].

There is a lot more to be said about the connection between power operations in elliptic cohomology and the notion of replicability in (generalized) Moonshine. We will come back to these topics at a different occasion.

²Compare e.g. [Teo03, (2.1)] with $b = 1$, $t = 1/z$ and $F(q) = g(z)$.

APPENDIX A. THOM CLASSES AND THE FIXED POINT FORMULA

A.1. Thom classes in equivariant cohomology theories.

Definition A.1. We say that a compatible family of equivariant cohomology theories $\{E_G\}$ has natural Thom classes for complex vector bundles, if for every complex G -vector bundle V over a pointed G -space X there exists a class $u_G^E(V) \in \tilde{E}^0(X^\vee)$ with the following properties:

(1) Naturality: If $f: X \rightarrow Y$ is a pointed G -map, then

$$u_G^E(f^*V) = f^*u_G^E(V).$$

(2) Multiplicativity: The family $\{E_G\}$ has external products, and if V is a complex G -vector bundle over X and W is a complex H -vector bundle over Y , then

$$u_{G \times H}^E(V \oplus W) = u_G^E(V) \boxtimes u_H^E(W).$$

(3) Periodicity: If V is a complex G -representation, then

$$u_G^E(V) \in \tilde{E}^0(\mathbb{S}^V)$$

is a unit in E^* .

(4) Change of groups: If $\alpha: H \rightarrow G$ is a map of groups, then

$$\text{res}_\alpha u_G^E(V) = u_H^E(V).$$

If G is the trivial group we omit it from the notation. We might also omit E from the notation if it is clear from the context which cohomology theory is meant.

Fix a group G .

Definition A.2. We say that E_G has natural Thom classes for complex vector bundles if it satisfies axioms (1) and (3) of the previous definition and

$$(2') \quad u_G(V \oplus W) = u_G(V) \otimes u_G(W).$$

Note that (2) and (4) together imply (2'). In the situation of Definition A.2, let

$$\alpha := u(\mathbb{C}) \in \tilde{E}^{-2}(\text{pt})$$

be the periodicity element, and

$$\tau_G(V) := u_G(V) \cdot \alpha^{-d} \in \tilde{E}^{2d}(X^\vee),$$

where $d = \dim_{\mathbb{C}} V$. Then the τ_G satisfy tom Dieck's axioms for equivariant Thom classes (c.f. [May96, p.335]). At the other hand, if E_G satisfies the axioms of [May96, p.335] and E^2 contains a unit, then the axioms of Definition A.2 follow.

Example A.3. Complex equivariant K -theory and Borel equivariant E theory for even periodic E satisfy the axioms of Definition A.1. For any E satisfying the axioms of [May96, p.335],

$$EP_G := \bigvee_{n \in \mathbb{Z}} \Sigma^{2n} E_G$$

satisfies the axioms of Definition A.2.

Recall (c.f. [Oko82], [May96]) that MU_G is universal among the G -equivariant cohomology theories with Thom classes in the sense of tom Dieck.

Proposition A.4. *The conditions of Definition A.2 are equivalent to the existence of a map of $\mathbf{G}$ -ring spectra*

$$\phi_{\mathbf{G}}: \mathrm{MUP}_{\mathbf{G}} \rightarrow \mathbf{E}_{\mathbf{G}}.$$

It is the unique map of ring spectra taking Thom classes to Thom classes.

PROOF : If $\phi_{\mathbf{G}}$ exists, we use it to push-forward the Thom classes of $\mathrm{MUP}_{\mathbf{G}}$ to $\mathbf{E}_{\mathbf{G}}$. In the situation of Definition A.2, the $\tau_{\mathbf{G}}$'s give rise to a map $\psi_{\mathbf{G}}: \mathrm{MU}_{\mathbf{G}} \rightarrow \mathbf{E}_{\mathbf{G}}$, and we set

$$\phi_{\mathbf{G}}|_{\Sigma^{2n}\mathrm{MU}_{\mathbf{G}}} := (\Sigma^{2n}\psi_{\mathbf{G}}) \cdot \alpha^n.$$

Then $\phi_{\mathbf{G}}$ is a map of $\mathbf{G}$ -ring spectra, and the two constructions are inverse to each other. $\square$
Traditionally, topologists like to work with graded rings and define the genus corresponding to $\psi_{\mathbf{G}}$ as the composite

$$\begin{aligned} \mathcal{N}_{\mathbf{u},\mathbf{G}}^* &\rightarrow \mathrm{MU}_{\mathbf{G}}^*(\mathrm{pt}) \rightarrow \mathbf{E}_{\mathbf{G}}^*(\mathrm{pt}) \\ [M] &\longmapsto \psi_{\mathbf{G}}(M), \end{aligned}$$

where the first map is the Pontrjagin-Thom map from the complex equivariant cobordism ring to the graded coefficient ring of $\mathrm{MU}_{\mathbf{G}}$. Alternatively, we can consider the composite of ungraded maps

$$\mathcal{N}_{\mathbf{u},\mathbf{G}}^* \rightarrow \mathrm{MUP}_{\mathbf{G}}^0(\mathrm{pt}) \rightarrow \mathbf{E}_{\mathbf{G}}^0(\mathrm{pt})$$

to define $\phi_{\mathbf{G}}(M)$. Hence if $[M] \in \mathcal{N}_{\mathbf{u},\mathbf{G}}^{2d}$, then

$$\psi_{\mathbf{G}}(M) = \phi_{\mathbf{G}}(M)\alpha^{-d}.$$

Example A.5. For the Atiyah-Bott-Shapiro K-theory Thom classes for complex vector bundles, $\mathbf{u}_{\mathbf{G}}^{\mathrm{Td}}$, the periodicity element α is the Bott element β , and we have $\phi_{\mathbf{G}}(M) = \mathrm{Td}(M)$ and $\psi_{\mathbf{G}}(M) = \mathrm{Td}_{\mathbf{G}}(M) \cdot \beta^{-d}$.

We write $\theta_{\mathbf{E}}$ for the Thom isomorphism

$$\theta_{\mathbf{E}}(x) = x \cdot \mathbf{u}_{\mathbf{E}}^{\mathbf{G}}.$$

Further, $z: X_+ \rightarrow X^V$ denotes the zero section, and

$$e_{\mathbf{G}}^{\mathbf{E}}(V) = z^*(\mathbf{u}_{\mathbf{G}}^{\mathbf{E}}(V)) \in \mathbf{E}_{\mathbf{G}}^0(X)$$

denotes the Euler class of V . Using the fact that Thom classes of trivial bundles are given by units, one extends the definitions of the $\mathbf{u}_{\mathbf{G}}^{\mathbf{E}}$ and $\theta_{\mathbf{E}}$ to virtual bundles in the usual way. Recall further that the push-forward along the map $\pi: M \rightarrow \mathrm{pt}$, where M is a stably almost complex oriented $\mathbf{G}$ -manifold, is defined as the composite

$$\pi_!: \mathbf{E}_{\mathbf{G}}^0(M) \rightarrow \widetilde{\mathbf{E}}_{\mathbf{G}}^0(M^{-\mathbf{T}M}) \rightarrow \widetilde{\mathbf{E}}_{\mathbf{G}}^0(\mathbb{S}^0),$$

where $\mathbf{T}M$ is the tangent bundle of M , the first map is the Thom isomorphism, and the second map is the Pontrjagin-Thom collapse.

The genus $\phi_{\mathbf{G}}$ is then computed as the push-forward of 1:

$$(14) \quad \phi_{\mathbf{G}}(M) = \pi_{\mathbf{G}!}(1) \in \widetilde{\mathbf{E}}_{\mathbf{G}}^0(\mathbb{S}^0).$$

Similarly, the renormalized genus $\psi_{\mathbf{G}}(M)$ is the graded push-forward of 1 which is obtained by using the graded Thom isomorphism associated to τ :

$$\psi_{\mathbf{G}}(M) = \pi_{\mathbf{G}!}^{\tau}(1) \in \widetilde{\mathbf{E}}^{-2d}(\mathbb{S}^0)$$

(cf. [Oko82], [CF66]).

A.1.1. *The $\hat{A}$ -genus and the Todd genus Thom classes.* For the remainder of this appendix, we will closely follow the exposition in [Mila]. Consider the inclusion

$$\mathbf{i}: \text{Spin}(\mathfrak{n}) \rightarrow \text{Spin}^{\mathbb{C}}(\mathfrak{n}) \cong (\text{Spin}(\mathfrak{n}) \times \mathbf{U}(1)) / (\mathbb{Z}/2),$$

and let $\mathbf{V}$ be a real, $8k$ -dimensional vector bundle over X which has a (chosen) $\text{Spin}(8k)$ -structure. Let

$$\mathbf{u}_{\text{ABS}}(\mathbf{V}) \in \text{KO}(X^{\mathbf{V}})$$

be the Atiyah-Bott-Shapiro Thom class of $\mathbf{V}$. Forgetting the Spin -structure and viewing $\mathbf{V}$ as a $\text{Spin}^{\mathbb{C}}(8k)$ -bundle via the forgetful map $\mathbf{i}$, we get a K -theory Thom class

$$\mathbf{u}_{\text{ABS}}^{\mathbb{C}}(\mathbf{V}) \in K(X^{\mathbf{V}}),$$

and a close look at the definition of these Thom classes on p.31 of [ABS64] shows:

$$\mathbf{u}_{\text{ABS}}^{\mathbb{C}}(\mathbf{V}) = \mathbf{u}_{\text{ABS}}(\mathbf{V}) \otimes_{\mathbb{R}} \mathbb{C}.$$

We further recall the map

$$\tilde{\mathbf{l}}: \mathbf{U}(k) \rightarrow \text{Spin}^{\mathbb{C}}(2k),$$

which allows us to view any complex vector bundle $\mathbf{V}$ as a $\text{Spin}^{\mathbb{C}}$ bundle. Atiyah, Bott and Shapiro prove [ABS64, Thm.11.6]

$$\mathbf{u}_{\text{Td}}(\mathbf{V}) = \mathbf{u}_{\text{ABS}}^{\mathbb{C}}(\mathbf{V}).$$

These statements are summarized in the following commuting diagram of maps of ring spectra:

$$\begin{array}{ccccc} \text{MSU} & \longrightarrow & \text{MSpin} & \xrightarrow{\hat{A}} & \text{KO} \\ \downarrow & & \downarrow & & \downarrow \otimes \mathbb{C} \\ \text{MU} & \longrightarrow & \text{MSpin}^{\mathbb{C}} & \longrightarrow & \text{K} \\ & \searrow & \text{Td} & \nearrow & \end{array}$$

Let $\mathbf{U}^2(\mathfrak{n})$ be the pull-back in the diagram

$$(15) \quad \begin{array}{ccc} \mathbf{U}^2(\mathfrak{n}) & \xrightarrow{j} & \text{Spin}(2\mathfrak{n}) \\ \pi \downarrow & & \downarrow \\ \mathbf{U}(\mathfrak{n}) & \longrightarrow & \text{SO}(2\mathfrak{n}). \end{array}$$

In particular, $\mathbf{U}^2(1)$ is the non-trivial double cover of $\mathbf{U}(1)$ and hence isomorphic to it. Then $\mathbf{U}^2(\mathfrak{n})$ is also the pull-back in the square

$$\begin{array}{ccc} \mathbf{U}^2(\mathfrak{n}) & \longrightarrow & \mathbf{U}^2(1) \\ \pi \downarrow & & \downarrow \\ \mathbf{U}(\mathfrak{n}) & \xrightarrow{\det} & \mathbf{U}(1) \end{array}$$

(cf. [Mila]). Vector bundles with a $\mathbf{U}^2$ -structure are bundles which have at the same time a complex structure and a Spin structure. The second pull-back square implies that these are exactly those complex bundles whose determinant bundle possess a square root, together with a choice of this square root.

Note that (15) gives rise to two different factorizations of the map

$$\mathbf{U}^2(\mathfrak{n}) \longrightarrow \mathrm{SO}(2\mathfrak{n})$$

through $\mathrm{Spin}^{\mathbb{C}}(2\mathfrak{n})$: The composition of j with the inclusion i becomes

$$(j, 1): \mathbf{U}^2(\mathfrak{n}) \rightarrow (\mathrm{Spin}(2\mathfrak{n}) \times \mathbf{U}(1))/(\mathbb{Z}/2).$$

while composing π with $\widetilde{l}$ yields the map

$$(j, \widetilde{\det}): \mathbf{U}^2(\mathfrak{n}) \rightarrow (\mathrm{Spin}(2\mathfrak{n}) \times \mathbf{U}(1))/(\mathbb{Z}/2)$$

Let $\mathbf{M}$ be a complex Clifford module. Then the two above maps make $\mathbf{M}$ into $\mathbf{U}^2$ representations, and if we denote the representation obtained from $(j, 1)$ by $\rho_{\mathbf{M}}$ then the other one becomes $\rho_{\mathbf{M}} \otimes_{\mathbb{C}} \widetilde{\det}$.

Let $\mathbf{P}$ be a principal $\mathbf{U}^2(\mathfrak{n})$ -bundle, and let

$$\mathbf{V} = \mathbf{P} \times_{\mathbf{U}^2(\mathfrak{n})} \mathbb{R}^{2\mathfrak{n}}$$

be its associated vector bundle. Viewing $\mathbf{V}$ as $\mathrm{Spin}^{\mathbb{C}}(2\mathfrak{n})$ bundle via the map $(j, 1)$ (i.e. by first viewing it as a $\mathrm{Spin}(2\mathfrak{n})$ bundle and then forgetting some of the structure), gives the complex Atiyah-Bott-Shapiro Thom class

$$\mathbf{u}_{\hat{\mathbf{A}}}^{\mathbb{C}}(\mathbf{V}) := \mathbf{u}_{\mathrm{ABS}}^{\mathbb{C}}(\mathbf{V}) = \chi_{\mathbf{V}}^{\mathbb{C}}(\mathbf{P} \times_{\mathbf{U}^2(\mathfrak{n})} \rho_{\mu^{\mathbb{C}}}).$$

Here $\chi_{\mathbf{V}}^{\mathbb{C}}$ and $\mu^{\mathbb{C}}$ are as defined in [ABS64] Note that this is really an abuse of notation, since $\mathbf{u}_{\mathrm{ABS}}^{\mathbb{C}}(\mathbf{V})$ depends on the $\mathrm{Spin}^{\mathbb{C}}$ structure of $\mathbf{V}$, not just on $\mathbf{V}$. Viewing $\mathbf{V}$ as $\mathrm{Spin}^{\mathbb{C}}$ -bundle via the other map $(j, \widetilde{\det})$, we obtain a different class for $\mathbf{u}_{\mathrm{ABS}}^{\mathbb{C}}(\mathbf{V})$, namely

$$(16) \quad \mathbf{u}_{\mathrm{Td}}(\mathbf{V}) = \mathbf{u}_{\mathrm{ABS}}^{\mathbb{C}}(\mathbf{V}) = \chi_{\mathbf{V}}^{\mathbb{C}}\left(\mathbf{P} \times_{\mathbf{U}^2(\mathfrak{n})} \left(\rho_{\mu^{\mathbb{C}}} \otimes_{\mathbb{C}} \widetilde{\det}\right)\right) = \mathbf{u}_{\hat{\mathbf{A}}}(\mathbf{V}) \otimes_{\mathbb{C}} \sqrt{\det(\mathbf{V})}.$$

(The associated fibers of these two bundles are the same as complex Clifford modules, but are viewed as $\mathbf{U}^2(\mathfrak{n})$ -representations in different ways. Hence the associated bundles are not necessarily isomorphic.) The class $\mathbf{u}_{\mathrm{Td}}(\mathbf{V})$ is the Todd genus Thom class of $\mathbf{V}$ viewed as a complex vector bundle. If $\mathfrak{n} = 4k$, we have

$$\mathbf{u}_{\hat{\mathbf{A}}}^{\mathbb{C}}(\mathbf{V}) = \mathbf{u}_{\mathrm{ABS}}(\mathbf{V}) \otimes_{\mathbb{R}} \mathbb{C},$$

where $\mathbf{u}_{\mathrm{ABS}}(\mathbf{V})$ is the KO-Thom class of $\mathbf{V}$ viewed as $\mathrm{Spin}(8k)$ -bundle.

A.2. Atiyah and Segal's Lefschetz fixed point formula. Let $\mathbf{E}$ be an even periodic cohomology theory. Fix a group $\mathbf{G}$ and an element $\mathfrak{g} \in \mathbf{G}$. Then

$$\mathbf{X} \mapsto \mathbf{E}^*(\mathbf{X}^{\mathfrak{g}}) =: \mathbf{F}_{\mathbf{G}}^*(\mathbf{X})$$

is a $\mathbf{G}$ -equivariant cohomology theory. Let $\mathbf{X}$ be a $\mathbf{G}$ -pace, and let $\mathbf{V}$ be a $\mathbf{G}$ -equivariant complex vector bundle over $\mathbf{X}$. Then the $\mathfrak{g}$ -fixed points of $\mathbf{V}$ form a vector bundle over $\mathbf{X}^{\mathfrak{g}}$, namely $\mathbf{V}_{\mathfrak{g}=1}$, the eigenbundle of the eigenvalue 1 of the action of $\mathfrak{g}$ on $\mathbf{V}|_{\mathbf{X}^{\mathfrak{g}}}$.

Assume that we have chosen a complex orientation of $\mathbf{E}$. Then $\mathbf{F}_{\mathbf{G}}$ inherits natural Thom classes

$$\mathbf{u}_{\mathbf{G}}^{\mathbf{F}}(\mathbf{V}) := \mathbf{u}^{\mathbf{E}}(\mathbf{V}_{\mathfrak{g}=1}) \in \widetilde{\mathbf{E}}^0((\mathbf{X}^{\mathfrak{g}})^{\mathfrak{g}}).$$

Let now $\mathbf{E}_{\mathbf{G}}$ be an equivariant version of $\mathbf{E}$ with natural Thom classes $\mathbf{u}_{\mathbf{G}}^{\mathbf{E}}$ continuing those of the complex orientation of $\mathbf{E}$, and assume that $\mathbf{E}_{(\mathfrak{g})}^0$ is flat over $\mathbf{E}^0$. Consider the natural transformation

$$\mathbf{r}: \mathbf{E}_{\mathbf{G}}(\mathbf{X}) \rightarrow \mathbf{E}_{(\mathfrak{g})}(\mathbf{X}^{\mathfrak{g}})$$

defined by

$$r := i^* \circ \text{res}|_{\langle g \rangle}^G,$$

where $i: X^g \rightarrow X$ is the inclusion of the fixed points.

Definition A.6. For a complex G -vector bundle V over X , we write $V_{g=\zeta}$ for the ζ -eigenbundle of the action of g on $V|_{X^g}$. Write $V_{g \neq 1}$ for the orthogonal complement of V_1 ,

$$V_{g \neq 1} = \bigoplus_{\zeta \neq 1} V_\zeta.$$

Example A.7. Let M be a smooth complex G -manifold, and let T its tangent bundle. Then $T_{g=1}$ is the tangent bundle TM^g of M^g , and $T_{g \neq 1}$ is the normal bundle N^g of i .

Proposition A.8. *For a complex G -vector bundle V over X , we have*

$$r(\theta_E(x)) = \theta_F(r(x) \cdot e_G^E(V_{g \neq 1})).$$

Its proof relies on the following lemma about relative zero sections:

Lemma A.9 ([Rud98]). *Let V and W be (equivariant) vector bundles over X , let*

$$s: W \rightarrow V \oplus W$$

be the inclusion, and write X^s for the induced map of Thom spaces

$$X^s: X^W \rightarrow X^{V \oplus W}.$$

Then X^s pulls back $u_G(V \oplus W)$ to

$$e_G(V) \cdot u_G(W) \in \tilde{E}_G^0(X^W).$$

PROOF OF PROPOSITION A.8: The fixed points inclusion

$$i: (X^V)^g \rightarrow X^V$$

factors into the composition of the two maps

$$i_1 := (X^g)^s: (X^g)^{V_{g=1}} \rightarrow (X^g)^{V_{g \neq 1} \oplus V_{g=1}}$$

and

$$i_2: (X^g)^{V_{g \neq 1} \oplus V_{g=1}} \rightarrow X^V,$$

where i_2 is the map of Thom spaces obtained from the fixed points inclusion $i_g: X^g \rightarrow X$ and the corresponding bundle map

$$\tilde{i}: i^*(V) \cong V_{g \neq 1} \oplus V_{g=1} \rightarrow V.$$

We have

$$\begin{aligned} r(\theta_E(x)) &= r(x \cdot u_G^E(V)) \\ &= r(x) \cdot r(u_G^E(V)) \\ &= r(x) \cdot i_1^* \circ i_2^* \text{res}|_{\langle g \rangle}^G u_G^E(V) \\ &= r(x) \cdot i_1^* \circ i_2^* u_{\langle g \rangle}^E(V) \\ &= r(x) \cdot i_1^* (u_{\langle g \rangle}^E(V_{g \neq 1} \oplus V_{g=1})) \\ &= r(x) \cdot e_{\langle g \rangle}^E(V_{g \neq 1}) \cdot u_{\langle g \rangle}^E(V_{g=1}) \\ &= \theta_F(r(x) \cdot e_{\langle g \rangle}^E(V_{g \neq 1})), \end{aligned}$$

where the second to last equality is Lemma A.9. Let α be the unique map from $\langle g \rangle$ to the trivial group. Since the action of $\langle g \rangle$ on $V_{g=1}$ is trivial, we have

$$u_{\langle g \rangle}^E(V_{g=1}) = \text{res}|_{\alpha} u^E(V_{g=1}) = u_G^F(V).$$

□

The correction factor $e_{\langle g \rangle}(V_{g \neq 1})$ is an exponential characteristic class. If we assume the Euler classes of $\langle g \rangle$ -representations to be invertible and set

$$e_{\langle g \rangle}(-V) := e_{\langle g \rangle}(V)^{-1} \in \widetilde{E}_{\langle g \rangle}^0(\mathbb{S}^V),$$

it follows that Proposition A.8 holds for virtual bundles, too. In the case $E_G = K_G$, and $u_G^E = u_G^{\text{Td}}$, the Riemann-Roch formula in [Dye69]

$$\pi_!^E(x) = \pi_!^F(r(x) \cdot e_{\langle g \rangle}^{\text{Td}}(-N^g))$$

becomes exactly the Atiyah-Segal result [AS68, 2.10]

$$(17) \quad (\text{ind}_G^X(x))_g = (\text{ind}_1^{X^g} \otimes \text{id}_{R(\langle g \rangle)})_g \left(\frac{r(x)}{\lambda_{-1}(N^g)} \right).$$

Here $\lambda_{-1}(V)$ denotes the alternating sum of the exterior powers of the $\langle g \rangle$ -representation V . Let now $u^{\hat{A}}$ be the complex Atiyah-Bott-Shapiro Thom class for $\text{Spin}(2d)$ -vector bundles. For simplicity, we assume that M is a U^2 -manifold. We combine the classical Riemann-Roch formula for the Chern character

$$\begin{aligned} \pi_!^{\hat{A}}(a) &= \pi_!^H \left(\frac{\text{ch}(e^{\hat{A}}(-TM))}{e^H(-TM)} \cdot \text{ch}(a) \right) \\ &= \int_M \hat{A}(TM) \text{ch}(a) \end{aligned}$$

with the Lefschetz formula (17) to obtain

$$\begin{aligned} (18) \quad \left({}^M \pi_{G!}^{\hat{A}}(a) \right)(g) &= \int_{M^g} \frac{e^H(TM^g)}{\text{ch}(e^{\hat{A}}(TM|_{M^g})(g))} \cdot \text{ch}((a|_{M^g})(g)) \\ &= \int_{M^g} \frac{e^H(TM^g) \text{ch}(\sqrt{\det TM|_{M^g}}(g))}{\text{ch}(e^{\text{Td}}(TM|_{M^g})(g))} \cdot \text{ch}((a|_{M^g})(g)) \\ &= \varepsilon \left(\left(\prod_{x_i} x_i \right) \left(\prod_{r=0}^{k-1} \prod_{y_i} \frac{e^{\frac{y_i}{2} + \pi i \frac{r}{k}}}{1 - e^{y_i + 2\pi i \frac{r}{k}}} \right) \cdot \text{ch}((a|_{M^g})(g)) \right) [M^g]. \end{aligned}$$

Here the x_i are the Chern roots of TM^g , for fixed r , the y_i run over the Chern roots of the $e^{2\pi i \frac{r}{k}}$ eigenbundle $(TM|_{M^g})$, and $\varepsilon = \pm 1$ depends on the way that the action of g on $\det TM$ lifts to an action on $\sqrt{\det TM}$ and on our choice of identification of $U^2(1)$ with $U(1)$ above.

Remark A.10. The above discussion remains valid if we replace $\langle g \rangle$ by a topologically cyclic group e.g. S^1 or $\mathbb{R}/\mathbb{L}\mathbb{Z}$.

Let now P be a principal $\text{Spin}(2n)$ bundle over X , and $V = P \times_{\text{Spin}(2n)} \mathbb{R}^{2n}$ its associated vector bundle. We assume that V has an even S^1 -action, i.e., one that is induced by an S^1 -action on P . We write

$$u_{S^1}^{\hat{A}}(V) := u_{S^1}^{\hat{A}, C}(V)$$

for its equivariant Atiyah-Bott-Shapiro Thom class in complex K-theory. Note that $V_{g=1}$ and its orthogonal complement are still even dimensional Spin bundles. Hence the above discussion goes through with $U(n)$ replaced by $Spin(2n)$, and if we let $\pi_{\dagger}^{\hat{A}}$ denote the (equivariant) push-forward in K-theory defined by $u^{\hat{A}}$, we get

$$\hat{A}\pi_{\mathbb{S}^1 \dagger}^X(x) = \hat{A}\pi_{\dagger}^{X^{\mathbb{S}^1}} \left(r(x) \cdot e_{\mathbb{S}^1}^{\hat{A}}(-N) \right),$$

where N is the normal bundle of the fixed point inclusion. In order to compute the equivariant Euler class of N (or more generally of $V_{g \neq 1}$), recall that, since $\mathbb{S}^1$ acts fibre preserving and fixed-point free on N , we have by (4)

$$(19) \quad N \cong_{\mathbb{R}} \bigoplus_{j \geq 1} V_j q^j,$$

and N can be equipped (uniquely) with a complex structure making this a complex isomorphism. Then (16) gives

$$(20) \quad \begin{aligned} e_{\mathbb{S}^1}^{\hat{A}}(N) &= e_{\mathbb{S}^1}^{\text{Td}}(N) \cdot \left(\sqrt{\det(N)} \wr \mathbb{S}^1 \right)^{-1} \\ &= q^{-d} \left(\sqrt{\det N} \right)^{-1} \prod_{j \geq 1} \wedge_{-q^j} V_j, \end{aligned}$$

where $d = \sum_j j \dim_{\mathbb{R}} V_j$.

REFERENCES

- [ABS64] M. F. Atiyah, R. Bott, and A. Shapiro. Clifford modules. *Topology*, 3(suppl. 1):3–38, 1964.
- [AFG] Matthew Ando, Christopher French, and Nora Ganter. The Jacobi orientation and the two-variable elliptic genus. Available at the arXiv as math.AT/0605554.
- [AHS01] M. Ando, M. J. Hopkins, and N. P. Strickland. Elliptic spectra, the Witten genus and the theorem of the cube. *Invent. Math.*, 146(3):595–687, 2001.
- [AHS04] Matthew Ando, Michael J. Hopkins, and Neil P. Strickland. The sigma orientation is an H_{∞} map. *Amer. J. Math.*, 126(2):247–334, 2004.
- [AM01] Matthew Ando and Jack Morava. A renormalized Riemann-Roch formula and the Thom isomorphism for the free loop space. In *Topology, geometry, and algebra: interactions and new directions (Stanford, CA, 1999)*, volume 279 of *Contemp. Math.*, pages 11–36. Amer. Math. Soc., Providence, RI, 2001.
- [And92] Matthew Ando. *Operations in complex-oriented cohomology theories related to subgroups of formal groups*. PhD thesis, Massachusetts Institute of Technology, 1992.
- [And00] Matthew Ando. Power operations in elliptic cohomology and representations of loop groups. *Trans. Amer. Math. Soc.*, 352(12):5619–5666, 2000.
- [AS68] M. F. Atiyah and G. B. Segal. The index of elliptic operators. II. *Ann. of Math. (2)*, 87:531–545, 1968.
- [BMMS86] R. R. Bruner, J. P. May, J. E. McClure, and M. Steinberger. H_{∞} ring spectra and their applications, volume 1176 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1986.
- [Bor92] Richard E. Borcherds. Monstrous moonshine and monstrous Lie superalgebras. *Invent. Math.*, 109(2):405–444, 1992.
- [CF66] P. E. Conner and E. E. Floyd. *The relation of cobordism to K-theories*. Lecture Notes in Mathematics, No. 28. Springer-Verlag, Berlin, 1966.
- [Dev96] Jorge A. Devoto. Equivariant elliptic homology and finite groups. *Michigan Math. J.*, 43(1):3–32, 1996.

- [dFLNU06] Tommaso de Fernex, Ernesto Lupercio, Thomas Nevins, and Bernardo Uribe. A localization principle for orbifold theories. In *Recent developments in algebraic topology*, volume 407 of *Contemp. Math.*, pages 113–133. Amer. Math. Soc., Providence, RI, 2006.
- [DLM02] Chongying Dong, Kefeng Liu, and Xiaonan Ma. On orbifold elliptic genus. In *Orbifolds in mathematics and physics (Madison, WI, 2001)*, volume 310 of *Contemp. Math.*, pages 87–105. Amer. Math. Soc., Providence, RI, 2002.
- [DMVV97] Robbert Dijkgraaf, Gregory Moore, Erik Verlinde, and Herman Verlinde. Elliptic genera of symmetric products and second quantized strings. *Comm. Math. Phys.*, 185(1):197–209, 1997.
- [Dye69] Eldon Dyer. *Cohomology theories*. Mathematics Lecture Note Series. W. A. Benjamin, Inc., New York-Amsterdam, 1969.
- [FQ93] Daniel S. Freed and Frank Quinn. Chern-Simons theory with finite gauge group. *Comm. Math. Phys.*, 156(3):435–472, 1993.
- [Gan] Nora Ganter. Twisted Hecke operators in equivariant elliptic cohomology and generalized Moonshine. arXiv:0706.2898.
- [Gan06] Nora Ganter. Orbifold genera, product formulas and power operations. *Advances in Mathematics*, 2006.
- [GKV] Victor Ginzburg, Mikhail Kapranov, and Eric Vasserot. Elliptic Algebras and Equivariant Elliptic Cohomology. arXiv:q-alg/9505012.
- [HBJ92] Friedrich Hirzebruch, Thomas Berger, and Rainer Jung. *Manifolds and modular forms*. Aspects of Mathematics, E20. Friedr. Vieweg & Sohn, Braunschweig, 1992. With appendices by Nils-Peter Skoruppa and by Paul Baum.
- [HKR00] Michael J. Hopkins, Nicholas J. Kuhn, and Douglas C. Ravenel. Generalized group characters and complex oriented cohomology theories. *J. Amer. Math. Soc.*, 13(3):553–594 (electronic), 2000.
- [KM] Nitu Kitchloo and Jack Morava. Thom Spectra for Loopgroup representations. arXiv:AT/0404541.
- [LUX] Ernesto Lupercio, Bernardo Uribe, and Miguel A. Xicoténcatl. The loop orbifold of the symmetric product. arXiv:math.AT/0606573.
- [May96] J. P. May. *Equivariant homotopy and cohomology theory*, volume 91 of *CBMS Regional Conference Series in Mathematics*. Published for the Conference Board of the Mathematical Sciences, Washington, DC, 1996. With contributions by M. Cole, G. Comezaña, S. Costenoble, A. D. Elmendorf, J. P. C. Greenlees, L. G. Lewis, Jr., R. J. Piacenza, G. Triantafyllou, and S. Waner.
- [MH02] Mark Mahowald and Mike Hopkins. The structure of 24 dimensional manifolds having normal bundles which lift to BO[8]. In *Recent progress in homotopy theory (Baltimore, MD, 2000)*, volume 293 of *Contemp. Math.*, pages 89–110. Amer. Math. Soc., Providence, RI, 2002.
- [Mila] Haynes Miller. The fixed point formula. Unpublished notes.
- [Milb] Haynes Miller. Loop spaces and the Witten genus. Unpublished notes.
- [Moe02] Ieke Moerdijk. Orbifolds as groupoids: an introduction. In *Orbifolds in mathematics and physics (Madison, WI, 2001)*, volume 310 of *Contemp. Math.*, pages 205–222. Amer. Math. Soc., Providence, RI, 2002.
- [Oko82] Christian Okonek. Der Conner-Floyd-Isomorphismus für Abelsche Gruppen. *Math. Z.*, 179(2):201–212, 1982.
- [Rud98] Yuli B. Rudyak. *On Thom spectra, orientability, and cobordism*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 1998. With a foreword by Haynes Miller.
- [Seg88] Graeme Segal. Elliptic cohomology (after Landweber-Stong, Ochanine, Witten, and others). *Astérisque*, (161-162):Exp. No. 695, 4, 187–201 (1989), 1988. Séminaire Bourbaki, Vol. 1987/88.
- [Tam] Hirotaka Tamasai. Infinite product decomposition of orbifold mapping spaces. Preprint, available at <http://arxiv.org/abs/0706.0932v1>.
- [tD68] Tammo tom Dieck. Steenrod-Operationen in Kobordismen-Theorien. *Math. Z.*, 107:380–401, 1968.
- [Teo03] Lee-Peng Teo. Analytic functions and integrable hierarchies—characterization of tau functions. *Lett. Math. Phys.*, 64(1):75–92, 2003.